arXiv:1111.0536v1 [math.PR] 2 Nov 2011

UNIQUENESS AND BLOW–UP FOR THE NOISY VISCOUS DYADIC MODEL

MARCO ROMITO

ABSTRACT. We consider the dyadic model with viscosity and additive Gaussian noise as a simplified version of the stochastic Navier–Stokes equations, with the purpose of studying uniqueness and emergence of singularities. We prove path–wise uniqueness and absence of blow–up in the intermediate intensity of the non–linearity, morally corresponding to the 3D case, and blow–up for stronger intensity. Moreover, blow–up happens with probability one for regular initial data.

1. INTRODUCTION

Motivations. Uniqueness is a problem with many facet for PDEs and different problems may require different approaches. When turning to stochastic PDEs, the problem acquires new levels of complexity, as uniqueness for stochastic processes can be understood in different ways (path–wise, in law, etc.). There are several recent result on this topic (see for instance [25] for a review).

A prototypical example of PDE without uniqueness are the Navier–Stokes equations, where the issue of uniqueness is mixed with the issue of regularity and emergence of singularities (see [22]). The analysis of the stochastic version of the equations has also received a lot of attention, and in recent years, by means of a clever way to solve the Kolmogorov equation, Da Prato and Debussche [13] (see also [19]) have shown existence of Markov families of solutions to the problem, proving that such Markov families admit, under suitable assumptions on the noise, a unique invariant measure, with exponential convergence rate [38]. In [26, 28] similar results have been obtained with a completely different method, based on the Krylov selection method [35]. Related results can be found in [27, 14, 24, 40, 43, 41, 44, 42, 1].

The two methods apply equally well in more general situations (as done for instance in [8]). The purpose of this paper is to analyse, with a view on uniqueness and emergence of blow–up, a much simpler infinite dimensional stochastic equation which anyway retains most of the characteristics of the original problem and which makes the methods of [13] and [28] applicable. To understand

---

Date: October 29, 2011.

2010 *Mathematics Subject Classification.* Primary 76D03; Secondary 35Q35 60H15 35R60.

Key words and phrases. dyadic model, infinite dimensional system of stochastic equations, path–wise uniqueness, blow–up.

which system could provide a good candidate, the essential point is to identify the non–linearity. If we look at the Navier–Stokes non–linearity on the torus with periodic boundary conditions, we have in Fourier variables,

$$(u\cdot\nabla)u = i\sum_{\mathbf{k}\in\mathbf{Z}^3,\mathbf{k}\neq 0}\Big(\sum_{\mathbf{l}+\mathbf{m}=\mathbf{k},\ \mathbf{l},\mathbf{m}\neq 0}(u_{\mathbf{l}}\cdot\mathbf{m})u_{\mathbf{m}}\Big)\,e^{i\mathbf{k}\cdot x},$$

that is, the $\mathbf{k}^{\text{th}}$ mode interacts with almost every other mode. The most reasonable simplification is to reduce the complexity of the interaction to a finite number of modes, while keeping the orthogonality property (which gives the conservation of energy), and the simplest is the nearest neighbour interaction. This is the dyadic model.

The dyadic model. The dyadic model has been introduced in [29, 33] as a model of the interaction of the energy of an inviscid fluid among different packets of wave–modes (shells). It has been lately studied in [34, 46, 12, 6, 4] and in the inviscid and stochastically forced case in [5, 9, 3]).

The viscous version has been studied in [30, 10, 11], and in particular [10] proves blow–up of *positive* solutions for the problem with non–linearity of strong intensity, and later in [7] the authors prove well–posedness and convergence to the inviscid limit, again for positive solutions, with non–linearity of intensity of “Navier–Stokes” type.

In this paper we study the dyadic model with additive noise,

$$dX_n = \big(-\nu\lambda_n^2 X_n + \lambda_{n-1}^\beta X_{n-1}^2 - \lambda_n^\beta X_n X_{n+1}\big)\,dt + \sigma_n\,dW_n, \qquad n\geqslant 1, \tag{1.1}$$

where $\lambda = 2$, $\lambda_n = \lambda^n$ and $X_0 \equiv 0$. The dispersion coefficients satisfy suitable assumptions (see Assumption 2.1) and the parameter β measures the relative intensity of the non–linearity with respect to the linear term.

The non–linear term *formally* satisfies the following property,

$$\sum_{n=1}^\infty X_n\big(\lambda_{n-1}X_{n-1}^2 - \lambda_n^\beta X_n X_{n+1}\big) = 0,$$

providing an a–priori bound of $X = (X_n)_{n\geqslant 1}$ in $\ell^2(\mathbf{R})$ which is independent of β. On the other hand the linear term and the non–linear term are roughly of the same size if

$$\lambda_n^2 X_n \approx \lambda_n^\beta X_n^2 \qquad \rightsquigarrow \qquad \lambda_n^{\beta-2}X_n \approx O(1),$$

which is the essential reason why local strong solutions exist when the initial condition decays at least as $\lambda_n^{-(\beta-2)}$ (see Theorem 2.4). If $\beta \leqslant 2$ this is always true due to the ℓ^2 bound explained above, and in fact [10] proves that the non–rndom problem has a unique global solution. The same method also works in the noisy case and it is straightforward to prove path–wise uniqueness in the case $\beta \leqslant 2$.

By a scaling argument (see for instance [10]), one can "morally" identify the dyadic model with the Navier–Stokes equations when $\beta \approx \frac{5}{2}$. In [7] well–posedness is proved in a range which includes the value $\frac{5}{2}$, but only for *positive* solutions (positivity is preserved by the unforced dynamics). It is clear that, as is, positivity is broken by the random perturbation.

Main results. This paper contains a thorough analysis of the case $\beta > 2$, which can be roughly summarised in the table below.

	$\beta \leqslant 2$	$2 < \beta \leqslant 3$	$\beta > 3$
blow–up	NO	NO*	YES
uniqueness	YES	YES	?

We prove path–wise uniqueness (Theorem 4.1) in the range $\beta \in (2, 3]$ by decomposing the solution in a quasi–positive component and a residual term. Quasi–positivity (see Section 3) is preserved by the system as long as the random perturbation is *not too strong* (the meaning of this will be explained below). Under the same conditions the residual term is small and provides a lower bound for the solution.

The quasi–positivity, together with the invariant area argument of [7], implies smoothness of the solution (Theorem 4.2), where by smoothness we mean that $(\lambda_n^\gamma X_n)_{n\geqslant 1}$ is bounded for every γ. This result holds for $\beta \in (2, \beta_c)$, where $\beta_c \in (2, 3]$ is the value idenfied in [7].

When $\beta > 3$ we use an idea of [10], which only works for positive solutions, together with quasi–positivity, to identify a set of initial conditions which leads to blow–up with positive probability (Theorem 6.1). While there are already cases where emergence of blow–up is proved, as for instance [16, 17] for the Schrödinger equation, [37, 36, 21] for the nonlinear heat equation (there is also [23], but their result is essentially one dimensional and no ideas for infinite dimensional systems are involved), such results essentially prove only that blow–up occurs with positive probability.

Our main result on blow–up for the dyadic model, Theorem 6.2, states that blow–up occurs with full probability, as long as the initial condition satisfies $\lambda_n^\alpha X_n(0) = O(1)$ for some $\alpha > \beta - 2$. This is optimal since it corresponds to the condition which ensures the existence of a local smooth solution, as remarked before. Essentially we prove that the $\lambda_n^{-(\beta-2)}$–decay is transient.

It remains open to understand uniqueness for $\beta > 3$, since blow–up rules out the use of smooth solutions, making path–wise uniqueness an harder problem. Uniqueness in law may still be achievable.

Methods. Our results are essentially based on the four following ideas, introduced in this paper to analyse the model.

Quasi–positivity. As already mentioned, the deterministic dynamics preserves positivity and an external forcing in principle destroys this. We prove that a negative lower bound can still be proved under the condition that the random perturbation is not too big (in a sense clarified in the next idea) and the negative part of the initial condition is small. This cannot be true for general initial states, and in fact we require this to be true only for the modes $X_n(0)$ of the initial condition corresponding to n large enough.

Irrelevance of the perturbation. The deterministic results of [10, 7] are not directly applicable due to the presence of the random term, responsible of the loss of positivity. Quasi–positivity works with a small perturbation and in general the random perturbation does not stay small.

We recover smallness in two ways, equally effective: on the one hand we do not need to have a small effect of the perturbation for every component, but only for large enough modes (in the duality frequency/wavelength this would correspond to small scales). On the other hand, due to viscosity, the effect of the randomness can be quantified in terms of regularity (i. e. decay in terms of powers of λ_n) and, roughly speaking, if the effect is finite with respect to some decay, then it is small with respect to any weaker decay.

Contraction of the negative components. In order to prove that blow–up happens with full probability, the above ideas are not sufficient, as they only insure that it happens with positive probability. We give a stronger form of quasi–positivity (Lemma 6.4), namely that the negative parts of the solution become smaller in a finite time, depending only on the size of the initial condition in H and on the size of the random perturbation.

Recurrence. We use the contraction of negative components to identify an event, spanning a time interval, thus an event for trajectories, that leads to a set where blow–up occurs. An argument of recurrence for these sets finally shows that blow–up has full probability. We remark that recurrence is not at all obvious, since for $\beta > 3$ the energy estimate is not strong enough to provide existence of a stationary solution with standard methods for dissipative stochastic PDEs.

Structure of the paper. Section 2 contains the basic definitions and assumptions, as well as the different notions of solution that will be used, together with some existence results. In Section 3 we show the control on negative components, which the basic result on which the paper is structured. Uniqueness and well–posedness are then proved in Section 4 for the intermediate range $2 < \beta \leqslant 3$. Section 5 contains some preliminary and general considerations on the blow–up time. These considerations are used in Section 6 to first identify events that lead to blow–up, then to prove that such sets are, conditional to the absence of blow–up, recurrent and hence that blow–up occurs with full probability.

Acknowledgements. The author wish to thank the *Institut Élie Cartan* of the *Université Henri Poincaré, Nancy I*, where part of this work has been done, for the kind hospitality and the exciting working environment. Warmful thanks to David Barbato, Franco Flandoli and Johnathan Mattingly for useful conversations and insights on the subject. This work is dedicated to Martina, she has arrived whilst the paper was being completed.

2. Preliminary results and definitions

We start by stating the assumptions on the strength of the noise we shall consider.

Assumption 2.1. The sequence $(\sigma_n)_{n\geqslant 1}$ of non–negative real numbers satisfies the following assumption of regularity: there is $\alpha_0 \in \mathbf{R}$ such that

$$\sup_{n\geqslant 1}\big(\lambda_n^{\alpha_0}\sigma_n\big) < \infty. \tag{2.1}$$

We shall see that, in order to ensure existence of solutions for problem (1.1), we need to impose a restriction on the possible values of α_0, depending on the value of the parameter β.

Assumption 2.2. If β is the parameter of problem (1.1), then the number α_0 of Assumption 2.1 above satisfies

$$\alpha_0 > \max\big\{\tfrac{1}{2}(\beta-3), \beta-3\big\} \tag{2.2}$$

2.1. **Notations.**

Set $\lambda = 2$ and $\lambda_n = \lambda^n$. For every $\alpha \in \mathbf{R}$ denote by V_α the (Hilbert) space

$$V_\alpha = \{(x_n)_{n\geqslant 1} : \sum_{n=1}^\infty (\lambda_n^\alpha x_n)^2 < \infty\},$$

with scalar product $\langle\cdot,\cdot\rangle_\alpha$ and norm $\|\cdot\|_\alpha$ given by

$$\langle x, y\rangle_\alpha = \sum_{n=1}^\infty \lambda_n^{2\alpha} x_n y_n, \qquad \|x\|_\alpha = \Big(\sum_{n=1}^\infty (\lambda_n^\alpha x_n)^2\Big).$$

Set in particular $H = V_0$ and $V = V_1$.

Let $\Omega_\beta = C([0,\infty); V_{-\beta})$ and define on Ω_β the canonical process $(\xi_t)_{t\geqslant 0}$ defined as $\xi_t(\omega) = \omega(t)$, $t \geqslant 0$. Define on Ω_β the canonical filtration $(\mathscr{B}_t)_{t\geqslant 0}$ where $\mathscr{B}_t$ is the Borel σ–field of $C([0,t]; V_{-\beta})$.

2.2. **Definitions of solution.**

We turn to the definition of solution. We shall consider first strong solutions, which are unique, regular but defined on a (possibly) random interval. Then we will consider weak solutions (in three different flavours).

2.2.1. *Strong solutions.* We first state the definition of local strong solution.

Definition 2.3 (Strong solution)**.** Let $\mathcal{W}$ be an Hilbert sub–space of H. Given a probability space $(\Omega, \mathscr{F}, \mathbb{P})$ and a cylindrical Wiener process $(W_t, \mathscr{F}_t)_{t\geqslant 0}$ on H, a *strong solution in* $\mathcal{W}$ with initial condition $x \in \mathcal{W}$ is a pair $(X(\cdot; x), \tau_x^{\mathcal{W}})$ such that

- $\tau_x^{\mathcal{W}}$ is a stopping time with $\mathbb{P}[\tau_x^{\mathcal{W}} > 0] = 1$,
- $X(\cdot; x)$ is a process defined on $[0, \tau_x^{\mathcal{W}})$ with $\widetilde{\mathbb{P}}[X(0, x) = x] = 1$,
- $X(\cdot; x)$ is continuous with values in $\mathcal{W}$ for $t < \tau_x^{\mathcal{W}}$,
- $\|X(t; x)\|_{\mathcal{W}} \to \infty$ as $t \uparrow \tau_x^{\mathcal{W}}$, $\widetilde{\mathbb{P}}$–a. s.,
- $X(\cdot; x)$ is solution of (1.1) on $[0, \tau_x^{\mathcal{W}})$.

The strong solution turns out to be a Markov process (and even a strong Markov process, but we do not need this fact here) in the following sense (see [31] for further details). Set $\mathcal{W}' = \mathcal{W} \cup \{\text{⚠}\}$, where the terminal state ⚠ is an isolated point. Define the set $\overline{W}(\mathcal{W}')$ of all paths $\omega : [0, \infty) \to \mathcal{W}'$ such that there exists a time $\zeta(\omega) \in [0, \infty]$ with ω continuous with values in $\mathcal{W}$ on $[0, \zeta(\omega))$ and $\omega(t) = \text{⚠}$ for $t \geqslant \zeta(\omega)$. The strong solution defined above can be extended as a process in $[0, \infty)$ with values in $\mathcal{W}'$ in a canonical way, achieving value ⚠ for $t \geqslant \tau_x^{\mathcal{W}}$. We say that the strong solution is Markov when the process on the extended state space $\mathcal{W}'$ is a Markov process.

Theorem 2.4. *Let* $\beta > 2$ *and assume* (2.1), (2.2). *Let* $\alpha \in (\beta - 2, \alpha_0 + 1)$, *then for every* $x \in V_\alpha$ *there exists a strong solution* $(X(\cdot; x), \tau_x^\alpha)$ *with initial condition* x.

Moreover, the solution is unique in the sense that if $(X(\cdot; x), \tau_x)$ *and* $(X'(\cdot; x), \tau_x')$ *are two solutions, then* $\mathbb{P}[\tau_x = \tau_x'] = 1$ *and* $X(\cdot; x) = X'(\cdot; x)$ *for* $t < \tau_x$.

Finally, the process $(X(\cdot; x))_{x \in V_\alpha}$ *is Markov, in the sense given above.*

Proof. Existence and uniqueness are essentially based on the same ideas of [42, Theorem 5.1], but with simpler estimates, we give a sketch of their proof because we will use some of the definitions we introduce later. Let $\chi \in C^\infty([0, \infty))$ be decreasing and such that $\chi(u) = 1$ for $u \leqslant 1$ and $\chi(u) = 0$ for $u \geqslant 2$, and consider the problem

$$dX_n^R = -\nu\lambda_n^2 X_n^R\, dt + \chi_R(\|X^R\|_\alpha)\big(\lambda_{n-1}^\beta (X_{n-1}^R)^2 - \lambda_n^\beta X_n^R X_{n+1}^R\big)\, dt + \sigma_n\, dW_n. \tag{2.3}$$

The problem has a (path–wise) unique global solution for every $x \in V_\alpha$, which is continuous in time with values in V_α. Given $x \in V_\alpha$, define $\tau_x^{\alpha,R}$ as the first time t when $\|X^R(t)\|_\alpha = R$, then $\tau_x^\alpha = \sup_{R>0} \tau_x^{\alpha,R}$ and the strong solution $X(t; x)$ coincides with $X^R(t; x)$ for $t \leqslant \tau_x^{\alpha,R}$. By uniqueness the definition makes sense. Markovianity follows by the Markovianity of each X^R. □

We notice that by path–wise uniqueness, if $x \in V_\alpha$, then $\tau_x^\alpha = \tau_x^{\alpha'}$ for every $\alpha' \in (\beta - 2, \alpha)$.

2.2.2. *Weak martingale solutions.* The fact that the *blow–up time* τ_x^α associated to a strong solution may (or may not) be infinite is the main topic of discussion of the paper. In order to have global solutions in the general case we state a weaker definition of solution.

Definition 2.5 (weak martingale solution)**.** Given $x \in H$, a probability measure $\mathbb{P}$ on Ω_β is a *weak martingale solution* of problem (1.1) with initial condition x if

1. for every $n \geqslant 1$, the process
$$M_t^n = \xi_n(t) - \xi_n(0) + \int_0^t \big(\nu\lambda_n^2 \xi_n(s) - \lambda_{n-1}^\beta \xi_{n-1}^2 + \lambda_n^\beta \xi_n \xi_{n+1}\big)\, ds,$$
defined through the canonical process ξ, is a continuous square integrable martingale with quadratic variation σ_n^2 (hence in particular a Brownian motion if $\sigma_n \neq 0$),
2. $\mathbb{P}[\xi(0) = x] = 1$.

It can be easily seen that this is equivalent to the standard definition of martingale solution (a proof is given for instance in [24] for the Navier–Stokes equations, but can be easily adapted to this problem).

Clearly both definition, of strong and weak solution, extend straightforwardly respectively to initial $\mathscr{F}_0$–measurable random variables and to initial distributions on H.

It turns out (see Theorem 2.9 below) that the strong solution represents the unique local solution, thus all weak solutions with the same initial condition coincide with it up to the *blow–up* time τ_x^α, if we require that weak solutions satisfy an additional property. Before stating the property, we need a few definitions for preparation.

Let $\mathbb{P}$ be a weak martingale solution and define for each n such that $\sigma_n \neq 0$ the process $W_n = \frac{1}{\sigma_n} M_n$ (and choose a W_n independent from all the other if $\sigma_n = 0$). Under $\mathbb{P}$ the $(W_n)_{n\geqslant 1}$ are independent standard one–dimensional Brownian motions and the sequence defined as

$$Z_n(t) = \sigma_n \Big(W_n(t) - \nu\lambda_n^2 \int_0^t e^{-\nu\lambda_n^2 (t-s)} W_n(s)\, ds \Big)$$

solves the following system,

$$\begin{cases} dZ_n + \nu\lambda_n^2 Z_n\, dt = \sigma_n\, dW_n, \\ Z_n(0) = 0, \end{cases} \qquad n \geqslant 1. \tag{2.4}$$

Moreover, if $Y = \xi - Z$, where ξ is the canonical process on Ω_β, thus a solution (in the sense of Definition 2.5) of (1.1) under $\mathbb{P}$, then $\mathbb{P}$–a. s.,

$$\dot{Y}_n + \nu\lambda_n^2 Y_n = \lambda_{n-1}^\beta (Y_{n-1} + Z_{n-1})^2 - \lambda_n^\beta (Y_n + Z_n)(Y_{n+1} + Z_{n+1}). \tag{2.5}$$

Define $\mathcal{G}_t$ as

$$\mathcal{G}_t(y,z) = \|y(t)\|_H^2 + 2\nu\int_0^t \|y(s)\|_V^2\,ds - \\ - 2\int_0^t \Big(\sum_{n=1}^\infty \lambda_n^\beta (y_n + z_n)(y_{n+1}z_n - y_n z_{n+1})\Big)\,ds,$$

it is easy to see (using the lemma below) that $\mathcal{G}_t(Y,Z)$ is finite if $Y \in L^\infty_{loc}(0,\infty;H)\cap L^2_{loc}(0,\infty;V)$ and Assumptions 2.1, 2.2 hold, and is jointly measurable in the variables (t,y,z) (see [8, 41] for a related problem). We first give a regularity result for Z, which is standard (see [15])

Lemma 2.6. *Assume* (2.1) *with* $\alpha_0 \in \mathbf{R}$. *Given* $\alpha < \alpha_0 + 1$, *then almost surely* $Z \in C([0,T];V_\alpha)$ *for every* $T > 0$. *Moreover, for every* $\epsilon \in (0,1]$, *with* $\epsilon < \alpha_0 + 1 - \alpha$, *there are* $c_{2.6\text{-}1.\epsilon} > 0$ *and* $c_{2.6\text{-}2.\epsilon} > 0$, *such that for every* $T > 0$,

$$\mathbb{E}\Big[\exp\Big(\frac{c_{2.6\text{-}2.\epsilon}}{T^\epsilon}\sup_{[0,T]}\|Z(t)\|_\alpha^2\Big)\Big] \leqslant c_{2.6\text{-}1.\epsilon}.$$

Definition 2.7 (energy martingale solution)**.** A weak martingale solution $\mathbb{P}$ with initial condition x is an *energy* martingale solution if

1. $\mathbb{P}[Y \in L^\infty_{loc}(0,\infty;H)\cap L^2_{loc}([0,\infty);V)] = 1$, where $Y = \xi - Z$ and Z is the solution to (2.4) associated to $\mathbb{P}$,
2. there is a set $T_{\mathbb{P}} \subset (0,\infty)$ of null Lebesgue measure such that for every $s \notin T_{\mathbb{P}}$ and every $t > s$,

$$\mathbb{P}[\mathcal{G}_t(Y,Z) \leqslant \mathcal{G}_s(Y,Z)] = 1.$$

Remark 2.8. Assume (2.1) with $\alpha_0 > 0$. As in [26, 27] one can consider an alternative definition where the almost sure energy inequality is replaced by the supermartingale condition given as follows:

- $\mathbb{P}[L^\infty(0,\infty;H)\cap L^2(0,\infty;V)] = 1$,
- for every $m \geqslant 1$ the process

$$\mathcal{E}_t^m = \|\xi(t)\|_H^{2m} + 2m\nu\int_0^t \|\xi(s)\|_H^{2m-2}\|\xi(s)\|_1^2\,ds - m(2m-1)\sigma^2\int_0^t \|\xi(s)\|_H^{2m-2}\,ds$$

  is an *almost sure supermartingale*, that is there is a set $T_{\mathbb{P}} \subset (0,\infty)$ of null Lebesgue measure such that for every $s \notin T_{\mathbb{P}}$ and every $t \geqslant s$,

$$\mathbb{E}^{\mathbb{P}}[\mathcal{E}_t^m|\mathscr{B}_s] \leqslant \mathcal{E}_s^m,$$

where $\sigma^2 = \sum_n \sigma_n^2$.

Given $\alpha > \beta - 2$ and $R > 0$, define the following random times on Ω_β,

$$\tau_\infty^\alpha = \inf\{t \geqslant 0 : \|\omega(t)\|_\alpha = \infty\}, \qquad \tau_\infty^{\alpha,R} = \inf\{t \geqslant 0 : \|\omega(t)\|_\alpha > R\}, \tag{2.6}$$

with the understanding that each random time is ∞ if the set is empty.

Theorem 2.9. *Let* $\beta > 2$ *and assume* (2.1), (2.2).

- *For every* $x \in H$ *there exists at least one energy martingale solution* $\mathbb{P}_x$.
- *If* $\alpha \in (\beta - 2, 1 + \alpha_0)$, $x \in V_\alpha$ *and* $\mathbb{P}_x$ *is an energy martingale solution with initial condition* x, *then* $\tau_x^\alpha = \tau_\infty^\alpha$ *under* $\mathbb{P}_x$ *and for every* $t > 0$,
$$\xi_s = X(s; x), \quad s \leqslant t, \qquad \mathbb{P}_x - a.s. \text{ on } \{\tau_x^\alpha > t\},$$
where $(X(\cdot; x), \tau_x^\alpha)$ *is the strong solution with initial condition* x *defined on* Ω_β.
- *There exists at least one family* $(\mathbb{P}_x)_{x \in H}$ *of energy martingale solutions satisfying the almost sure Markov property, namely for every* $x \in H$ *and every bounded measurable* $\phi : H \to \mathbf{R}$,
$$\mathbb{E}^{\mathbb{P}_x}[\phi(\xi_t)|\mathscr{B}_s] = \mathbb{E}^{\mathbb{P}_{\omega(s)}}[\phi(\xi_{t-s})], \qquad \mathbb{P}_x - a.s.,$$
for almost every $s \geqslant 0$ *(including* 0*) and for all* $t \geqslant s$.

Proof. The proof of the first fact can be done as in [2]. The proofs of the other two facts are entirely similar to those of Theorem 2.1 of [41] and Theorem 3.6 of [42] and we refer to these reference for further details. □

A natural way to prove existence of weak solution (and in fact this is the way it is done in [2]) is to use finite dimensional approximations, namely, consider for each $N \geqslant 1$ the solution $(X_n^{(N)})_{1 \leqslant n \leqslant N}$ to the following finite dimensional system,

$$\begin{cases} \dot{X}_1^{(N)} = -\nu\lambda_1^2 X_1^{(N)} - \lambda_1^\beta X_1^{(N)} X_2^{(N)} + \sigma_1 \, dW_1, \\ \dots, \\ \dot{X}_n^{(N)} = -\nu\lambda_n^2 X_n^{(N)} + \lambda_{n-1}^\beta (X_{n-1}^{(N)})^2 - \lambda_n^\beta X_n^{(N)} X_{n+1}^{(N)} + \sigma_n \, dW_n, \\ \dots, \\ \dot{X}_N^{(N)} = -\nu\lambda_N^2 X_N^{(N)} + \lambda_{N-1}^\beta (X_{N-1}^{(N)})^2 + \sigma_N \, dW_N, \end{cases} \tag{2.7}$$

Given $x \in H$, let $\mathbb{P}_x^{(N)}$ be the probability distribution on Ω_β of the solution of the above system with initial condition $x^{(N)} = (x_1, x_2, \dots, x_N)$.

Definition 2.10 (Galerkin martingale solution)**.** Given $x \in H$, a *Galerkin* martingale solution is any limit point in Ω_β of the sequence $(\mathbb{P}_x^{(N)})_{N \geqslant 1}$.

It is easy to verify (it is indeed the proof of existence in Theorem 2.9 and we refer to [2] for details in a similar problem) that *Galerkin* martingale solutions are *energy* martingale solutions.

Remark 2.11. The results stated in this section are essentially independent of the structure of the non–linear part and are only based on the fact that that the non–linearity is polynomial and at each mode involves only a finite number of other modes.

The results of the rest of the paper are on the other hand strongly based on the structure of the non–linear part. Besides the physical motivations of [29], from the point of view of studying a simplified version of Navier–Stokes, there are several possibilities for *nearest–neighbour* interaction among modes. One can write [34] any nearest–neighbour non–linearity as

$$\dot{X}_n = -\nu\lambda_n^2 X_n + (a_1 B_n^1(X) + a_2 B_n^2(X)),$$

where B_n^1 is the one corresponding to the dyadic model and $B_n^2(x) = \lambda_{n+1}^\beta x_{n+1}^2 - \lambda_n^\beta x_{n-1} x_n$. In [34] the authors notice that the inviscid problem with non–linearity B_n^2 is well–posed. Hence the dyadic model is the difficult, hence the most meaningful, part of any nearest–neighbour non–linearity.

3. CONTROL OF THE NEGATIVE COMPONENTS

Given $\beta > 2$, $\alpha \in \mathbf{R}$ and $c_0 > 0$, consider the solution Z of (2.4) and define the following process,

$$N_{\alpha,c_0}(t) = \min\big\{m \geqslant 1 : |Z_n(s)| \leqslant c_0 \nu \lambda_{n-1}^{-\alpha} \text{ for } s \in [0,t] \text{ and } n \geqslant m\big\}, \tag{3.1}$$

with $N_{\alpha,c_0}(t) = \infty$ if the set is empty.

Lemma 3.1 (Moments of N_{α,c_0}). *Given $\beta > 2$, assume (2.1) and let $\alpha < \alpha_0 + 1$. Then for every $\gamma \in (0, \alpha_0 + 1 - \alpha)$ and $\epsilon \in (0,1]$, with $\epsilon < \alpha_0 + 1 - \alpha - \gamma$, there are two numbers $c_{3.1\text{-}1} > 0$ and $c_{3.1\text{-}2} > 0$, depending only on ϵ, γ and α_0, such that for every $t > 0$ and $n \geqslant 1$,*

$$\mathbb{P}[N_{\alpha,c_0}(t) > n] \leqslant c_{3.1\text{-}1}\, e^{-c_{3.1\text{-}2} \frac{c_0 \nu}{t^\epsilon} \lambda_n^\gamma}.$$

In particular,

$$\mathbb{E}\Big[e^{\lambda^\gamma_{N_{\alpha,c_0}(t)}}\Big] < \infty,$$

and $\mathbb{P}[N_{\alpha,c_0}(t) = n] > 0$ for every $n \geqslant 1$.

Proof. For $n \geqslant 1$,

$$\{N_{\alpha,c_0}(t) \leqslant n\} = \Big\{\sup_{k \geqslant n} \sup_{[0,t]} \lambda_{k-1}^\alpha |Z_k(s)| \leqslant c_0 \nu\Big\},$$

hence if $\gamma < \alpha_0 + 1 - \alpha$ and $k \geqslant n$,

$$\sup_{[0,t]} \lambda_{k-1}^\alpha |Z_k(s)| \leqslant \lambda_{n-1}^{-\gamma} \sup_{[0,t]} \|Z(s)\|_{\alpha+\gamma},$$

and therefore by Chebychev's inequality and Lemma 2.6,

$$\mathbb{P}[N_{\alpha,c_0}(t) > n] \leqslant \mathbb{P}\Big[\sup_{[0,t]} \|Z(s)\|_{\alpha+\gamma} > c_0 \nu \lambda_{n-1}^\gamma\Big] \leqslant c_{2.6\text{-}1.\epsilon}\, e^{-c_{2.6\text{-}2.\epsilon} \frac{c_0 \nu}{t^\epsilon} \lambda_{n-1}^\gamma},$$

for every $\epsilon \in (0,1]$ with $\epsilon < \alpha_0 + 1 - \alpha - \gamma$. The double–exponential moment follows easily from this estimate.

We finally prove that $\mathbb{P}[N_{\alpha,c_0}(t)=n]>0$. We prove it for $n=1$, all other cases follow similarly. By independence,

$$\mathbb{P}[N_{\alpha,c_0}(t)=1]=\exp\Big(-\sum_{n=1}^{\infty}-\log\mathbb{P}\big[\sup_{[0,t]}\lambda_{k-1}^{\alpha}|Z_k(s)|\leqslant c_0\nu\big]\Big)$$

and it is sufficient to show that the series above is convergent. By (2.1),

$$\mathbb{P}\big[\sup_{[0,t]}\lambda_{k-1}^{\alpha}|Z_k(s)|\leqslant c_0\nu\big]\geqslant\mathbb{P}\big[\sup_{[0,t]}|\zeta(\lambda_n^2 s)|\leqslant 2^{\alpha}c_0\nu\lambda_n^{\alpha_0+1-\alpha}\big],$$

where ζ is the solution of the one–dimensional SDE $d\zeta+\nu\zeta\,dt=dW$, with $\zeta(0)=0$, and the conclusion follows by the fact that $\alpha<1+\alpha_0$ and standard tail estimates on the one–dimensional Ornstein–Uhlenbeck process (see for instance [20]). □

The lemma below is the crucial result of the paper. To formulate its statement, we introduce the finite dimensional approximations of the problem. Consider problem (2.5) and, for an integer $N\geqslant 1$, the finite dimensional approximations of (2.5),

$$\begin{cases}\dot Y_1^{(N)}=-\nu\lambda_1^2Y_1^{(N)}-\lambda_1^{\beta}X_1^{(N)}X_2^{(N)},\\ \dots,\\ \dot Y_n^{(N)}=-\nu\lambda_n^2Y_n^{(N)}+\lambda_{n-1}^{\beta}(X_{n-1}^{(N)})^2-\lambda_n^{\beta}X_n^{(N)}X_{n+1}^{(N)},\\ \dots,\\ \dot Y_N^{(N)}=-\nu\lambda_N^2Y_N^{(N)}+\lambda_{N-1}^{\beta}(X_{N-1}^{(N)})^2,\end{cases}\tag{3.2}$$

where for $n=1,\dots,N$ we have set $X_n^{(N)}=Y_n^{(N)}+Z_n$. It is easy to verify that the above SDE admits a unique global solution.

Lemma 3.2 (Main lemma). *Let $\beta>2$ and assume (2.1), (2.2). Let $\alpha\in[\beta-2,1+\alpha_0)$ and consider $C_0>0$, $a_0>0$ and $n_0\geqslant 1$ such that*

$$c_0\leqslant a_0\qquad\text{and}\qquad c_0<\sqrt{a_0}\big(\lambda_{n_0}^{\frac12(\alpha+2-\beta)}-\sqrt{a_0}\big).\tag{3.3}$$

Given $T>0$, let $N\geqslant 1$ and assume that $\lambda_{n-1}^{\alpha}X_n^{(N)}(0)\geqslant-a_0\nu$ for all $n=n_0,\dots,N$.

If $N>N_{\alpha,c_0}(T)$, then $Y_n^{(N)}(t)\geqslant-a_0\nu\lambda_{n-1}^{-\alpha}$ for all $t\in[0,T]$ and all $n\geqslant n_0\vee N_{\alpha,c_0}(T)$.

Proof. For simplicity we drop in this proof the superscript $^{(N)}$. We can first assume that $\lambda_{n-1}^{\alpha}Y_n(0)>-\nu a_0$ for $n\geqslant n_0,\dots,N$ (the equality will be included by continuity), then the same is true in a neighbourhood of $t=0$. Let $t_0>0$ be the first time when at least for one n, $\lambda_{n-1}^{\alpha}Y_n(t_0)=-\nu a_0$. Let $n\geqslant n_0\vee N_{\alpha,c_0}(T)$

be one of such indices, then

$$\begin{aligned}\dot{Y}_n(t_0) &\geqslant -\nu\lambda_n^2 Y_n(t_0) - \lambda_n^\beta(Y_n(t_0)+Z_n(t_0))(Y_{n+1}(t_0)+Z_{n+1}(t_0))\\ &\geqslant a_0\nu^2\lambda^2\lambda_{n-1}^{2-\alpha} + \lambda_n^\beta\big(a_0\nu\lambda_{n-1}^{-\alpha} - Z_n(t_0)\big)(Y_{n+1}(t_0)+Z_{n+1}(t_0))\\ &\geqslant a_0\nu^2\lambda^2\lambda_{n-1}^{2-\alpha} - \lambda_n^\beta\big(a_0\nu\lambda_{n-1}^{-\alpha} - Z_n(t_0)\big)(Y_{n+1}(t_0)+Z_{n+1}(t_0))_-\end{aligned}$$

since $\nu a_0\lambda_{n-1}^{-\alpha} - Z_n(t_0) \geqslant 0$ for $n \geqslant N_{\alpha,c_0}(T)$, and where $x_- = \max(-x,0)$. We also know that $Y_{n+1}(t_0) \geqslant -a_0\nu\lambda_n^{-\alpha}$, hence $Y_{n+1}(t_0)+Z_{n+1}(t_0) \geqslant -\nu(a_0+c_0)\lambda_n^{-\alpha}$ and so $(Y_{n+1}(t_0)+Z_{n+1}(t_0))_- \leqslant \nu(a_0+c_0)\lambda_n^{-\alpha}$. We also have $a_0\nu\lambda_{n-1}^{-\alpha} - Z_n(t_0) \leqslant \nu(a_0+c_0)\lambda_{n-1}^{-\alpha}$, so in conclusion

$$\dot{Y}_n(t_0) \geqslant \nu^2\lambda^2\lambda_{n-1}^{2-\alpha}\big(a_0 - \lambda_n^{\beta-2-\alpha}(a_0+c_0)^2\big) > 0$$

and the lemma is proved. □

The next theorem is based on Lemma 3.2 and shows that the process can explode *only* in the positive area.

Theorem 3.3. *Given* $\beta > 2$, *assume* (2.1) *and* (2.2). *Let* $\alpha \in (\beta-2, \alpha_0+1)$ *and* $x \in V_\alpha$, *and let* $(X(\cdot;x),\tau_x^\alpha)$ *be the strong solution in* V_α *with initial condition* x. *Then for every* $T > 0$ *and* $p \geqslant 1$,

$$\mathbb{E}\Big[\sup_{n\geqslant 1}\sup_{t\in[0,T\wedge\tau_x^\alpha]}\big(\lambda_{n-1}^\alpha(X_n(t))_-\big)^p\Big] < \infty.$$

In particular,

$$\inf_{n\geqslant 1}\inf_{t\in[0,\tau_x^\alpha\wedge T]}\lambda_{n-1}^\alpha X_n > -\infty, \qquad \mathbb{P}\text{–a. s.}$$

Proof. Fix $x \in V_\alpha$ and $T > 0$, and set $a_0 = \frac{1}{4}$ and $c_0 = \frac{1}{6}$, so that condition (3.3) holds for any n_0. Next, choose $n_0 \geqslant 1$ as the smallest integer such that $\lambda_{n-1}^\alpha x_n \geqslant -\frac{1}{4}\nu$ for all $n \geqslant n_0$. With the choice $c_0 = \frac{1}{6}$, set $\mathcal{Z}_{\alpha,T} = \{N_{\alpha,\frac{1}{6}}(T) < \infty\}$, which by Lemma 3.1 is an event of probability one.

Lemma 3.2 implies that on $\{\tau_x^\alpha > T\}$,

$$Y_n(t) \geqslant -\frac{1}{4}\nu\lambda_{n-1}^{-\alpha}, \qquad \text{for } n \geqslant n_0 \vee N_{\alpha,\frac{1}{6}}(T).$$

Indeed, we can set $x^{(N)} = (x_1,\dots,x_N)$ and notice that on the event $\{\tau_x^\alpha > T\}$, problem (2.5) has a unique solution, hence for every N the solution of (3.2) with initial condition $x^{(N)}$ converges to the solution of (2.5) with initial condition x (where the convergence is component–wise uniform in time on $[0,T]$).

Let $N_1 = n_0 \vee N_{\alpha,\frac{1}{6}}(T)$, it is clear that N_1 has the same finite moments of $N_{\alpha,\frac{1}{6}}(T)$, moreover on $\{\tau_x^\alpha > T\}$,

$$\lambda_{n-1}^\alpha X_n(t) \geqslant \begin{cases} -\lambda_{N_1-1}^\alpha \sup_{t\in[0,T]}\|X(t)\|_H, & n < N_1,\\ -\frac{5}{12}\nu & n \geqslant N_1,\end{cases}$$

for every $n \geqslant 1$, and so

$$\sup_{n\geqslant 1}\sup_{t\in[0,T]}\lambda_{n-1}^{\alpha}\big(X_n(t)\big)_{-} \leqslant \frac{5}{12}\nu + \lambda_{N_1-1}^{\alpha}\sup_{t\in[0,T]}\|X(t)\|_H$$

From Lemma 3.1 and the fact that $\mathbb{E}[\sup_{[0,T]}\|X(t)\|_H^p]$ is finite for every $p \geqslant 1$, the estimate in the statement of the theorem readily follows. $\square$

Remark 3.4. Given an initial condition $x \in V_\alpha$, if we set

$$\tau_{x,\pm}^{\alpha} = \sup\{t : \sup_{n\geqslant 1}\lambda_n^{\alpha}(X_n)_{\pm} < \infty\},$$

then $\tau_x^\alpha = \min(\tau_{x,+}^\alpha, \tau_{x,-}^\alpha)$, and the previous theorem essentially states that $\tau_x^\alpha = \tau_{x,+}^\alpha$.

Corollary 3.5. *Given* $\beta > 2$, *assume* (2.1) *and* (2.2). *Given* $\alpha \in (\beta - 2, \alpha_0 + 1)$ *and* $x \in V_\alpha$, *assume additionally either that problem* (2.5), *with initial condition* x, *admits a unique solution, for almost every possible value assumed by* Z, *or that we are dealing with a* Galerkin *solution starting in* x. *Then for every* $T > 0$ *and* $p \geqslant 1$,

$$\mathbb{E}\Big[\sup_{n\geqslant 1}\sup_{t\in[0,T]}\big(\lambda_{n-1}^{\alpha}\big(X_n(t)\big)_{-}\big)^p\Big] < \infty.$$

In particular,

$$\inf_{n\geqslant 1}\inf_{t\in[0,T]}\lambda_{n-1}^{\alpha}X_n > -\infty, \qquad \mathbb{P}\text{–a. s.}$$

Proof. We simply notice that in the proof of the theorem above we have used the piece of information $\tau_x^\alpha > T$ only to ensure that problem (2.5) admits a unique solution.

On the other hand, if we are dealing with a Galerkin solution, then up to a sub–sequence we still have component–wise uniform convergence in time. $\square$

4. UNIQUENESS AND REGULARITY FOR $2 < \beta \leqslant \frac{5}{2}$

In this section we prove two results, the first concerning path–wise uniqueness, the second concerning regularity (absence of blow–up), which are essentially extensions of the corresponding results for the case without noise. The extension is made possible by means of the control of negative components shown in Section 3. The path–wise uniqueness result below is restricted to *Galerkin* solutions (see Definition 2.10).

Theorem 4.1 (Path–wise uniqueness). *Let* $\beta \in [2, 3]$ *and assume that* (2.1), (2.2) *hold. Let* $X(0) \in V_{\beta-2}$, *then there exists a (path–wise) unique solution of* (1.1) *with initial condition* $X(0)$, *in the class of* Galerkin *martingale solutions.*

We do not know if there is uniqueness in some larger class (energy martingale solutions or weak martingale solutions), neither we know if a Galerkin martingale solution may develop blow–up. By slightly restricting the range of values of β, we do actually have an improvement over the previous result.

Theorem 4.2 (Smoothness). *There exists* $\beta_c \in (\frac{5}{2}, 3]$ *such that the following statement holds true. Assume that* (2.1), (2.2) *hold and let* $\beta \in (2, \beta_c)$ *and* $\alpha \in (\beta - 2, 1 + \alpha_0)$, *then for every* $x \in V_\alpha$, $\tau_x^\alpha = \infty$. *In particular, path–wise uniqueness holds in the class of* energy *martingale solutions.*

4.1. **The proof of Theorem 4.1.** The proof is based on the idea in [7, Proposition 3.2] which builds up on a result in [4]. Both results are proved for positive solutions (and no noise).

Proof of Theorem 4.1. It is sufficient to show uniqueness on any finite time interval, so we fix an arbitrary $T > 0$ and show that there is only one solution on $[0, T]$. We will use Lemma 3.2 with $c_0 = \frac{1}{6}$ and $a_0 = \frac{1}{4}$ (so that (3.3) holds for any n_0). In fact, since a Galerkin martingale solution is the component–wise limit of finite dimensional approximations, the bounds of the lemma remain stable in the limit to the infinite dimensional system.

Since $X(0) \in V_{\beta-2}$, we know that there is $n_0 \geqslant 1$ such that $\lambda_n^{\beta-2} X_n(0) \geqslant -\frac{1}{4}\nu$. We set $N_0 = 1 + n_0 \vee N_{\beta-2,\frac{1}{6}}(T)$.

Let X^1, X^2 two solutions with the same given initial condition $X^1(0) = X^2(0) = X(0)$. By Lemma 3.2 we know that $X_n^i(t) \geqslant Z_n(t) - \frac{1}{4}\nu\lambda_{n-1}^{2-\beta}$ for $n \geqslant N_0$, $t \in [0, T]$ and $i = 1, 2$. Set now

$$A_n = X_n^1 - X_n^2, \qquad B_n = \frac{1}{2}\nu\lambda_{n-1}^{2-\beta} - 2Z_n$$

and

$$\psi_\ell(t) = \sum_{n=1}^{N_0-1} \frac{A_n^2}{\lambda_n}, \qquad \psi_{h,N}(t) = \sum_{N_0}^{N} \frac{A_n^2}{\lambda_n}, \qquad \psi_N(t) = \psi_\ell(t) + \psi_{h,N}(t),$$

and notice that $X_n^1 + X_n^1 + B_n \geqslant 0$ if $t \in [0, T]$ and $n \geqslant N_0$. A simple computation shows that

$$\begin{aligned}\dot{A}_n = -\nu\lambda_n^2 A_n + \lambda_{n-1}^\beta (X_{n-1}^1 + X_{n-1}^2) A_{n-1} + \\ - \frac{1}{2}\lambda_n^\beta \big[(X_n^1 + X_n^2) A_{n+1} + (X_{n+1}^1 + X_{n+1}^2) A_n\big],\end{aligned}$$

hence for $N > N_0$,

$$\begin{aligned}\frac{d}{dt}\psi_{h,N} + 2\nu \sum_{n=N_0}^{N} \lambda_n A_n^2 = -\sum_{n=N_0}^{N} \lambda_n^{\beta-1}(X_{n+1}^1 + X_{n+1}^2) A_n^2 + \\ - \lambda_N^{\beta-1}(X_N^1 + X_N^2) A_N A_{N+1} + \lambda_{N_0-1}^{\beta-1}(X_{N_0-1}^1 + X_{N_0-1}^2) A_{N_0-1} A_{N_0} = \\ = \boxed{1} + \boxed{2}_N + \boxed{3}_{N_0}.\end{aligned}$$

For the first term we notice that $B_{n+1} \leqslant \frac{5}{6}\nu\lambda_n^{2-\beta}$, hence

$$\boxed{1} \leqslant \sum_{n=N_0}^{N} \lambda_n^{\beta-1} B_{n+1} A_n^2 \leqslant \nu \sum_{n=N_0}^{N} \lambda_n A_n^2.$$

For the second term,

$$\sum_{N=1}^{\infty} \int_0^T \boxed{2}_N \, dt \leqslant \sup_{[0,T]} \|X^1 + X^2\|_H \int_0^T \|A\|^2_{\frac{1}{2}(\beta-1)} \, ds,$$

and the quantity on the right–hand side is a. s. finite since by (2.1) Z is a. s. in $C([0,T]; V_\gamma)$ for every $\gamma < 1 + \alpha_0$ (hence for $\gamma = \frac{1}{2}(\beta - 1)$), and since by Definition 2.7, $V \in L^2([0,T]; V)$ (and $\frac{1}{2}(\beta-1) \leqslant 1$). This implies that a. s. $\int_0^T \boxed{2}_N \, dt \to 0$ as $N \to \infty$.

Likewise (recall that $X_0^1 = X_0^2 = 0$),

$$\frac{d}{dt}\psi_\ell \leqslant -\sum_{n=1}^{N_0-1} \lambda_n^{\beta-1}(X^1_{n+1} + X^2_{n+1})A_n^2 - \boxed{3}_{N_0} \leqslant \lambda^\beta_{N_0-1}\big(\sup_{[0,T]} \|X^1 + X^2\|_H\big)\psi_\ell - \boxed{3}_{N_0},$$

and in conclusion

$$\frac{d}{dt}\psi_N \leqslant \lambda^\beta_{N_0-1}\big(\sup_{[0,T]} \|X^1 + X^2\|_H\big)\psi_\ell + \boxed{2}_N.$$

Integrate in time (recall that $\psi_N(0) = 0$) and take the limit as $N \uparrow \infty$,

$$\psi(t) \leqslant \lambda^\beta_{N_0-1}\big(\sup_{[0,T]} \|X^1 + X^2\|_H\big) \int_0^t \psi(s) \, ds$$

where $\psi_N \uparrow \psi$ and $\psi(t) = \|A(t)\|^2_{-\frac{1}{2}}$. Since the term $\lambda^\beta_{N_0-1}\big(\sup_{[0,T]} \|X^1 + X^2\|_H\big)$ is a. s. finite, by Gronwall's lemma it follows that a. s. $A(t) = 0$ for all $t \in [0,T]$. ☐

4.2. **The proof of Theorem 4.2.** We start by giving a minimal requirement for smoothness of solutions to (1.1) which is analogous to the criterion developed in the noise–less case (see [7]).

Given $T > 0$ define the subspace K_T of Ω_β as

$$K_T = \big\{\omega \in \Omega_\beta : \lim_n \big(\max_{t \in [0,T]} \lambda_n^{\beta-2}|\omega_n(t)|\big) = 0\big\}.$$

Clearly $L^\infty(0,T; V_\alpha) \subset K_T$ for all $\alpha > \beta - 2$.

Proposition 4.3. *Under the assumptions of Theorem 4.2, let $x \in V_\alpha$. If $\mathbb{P}_x$ is an energy martingale solution (Definition 2.7) with initial condition x and τ_∞^α is the random time defined in (2.6), then for every $T > 0$,*

$$\{\tau_\infty^\alpha > T\} = K_T, \qquad \textit{under } \mathbb{P}_x.$$

Proof. Fix $\alpha \in (\beta - 2, \alpha_0 + 1)$, $x \in V_\alpha$ and an energy martingale solution $\mathbb{P}_x$ starting at x, and let τ_∞^α be the random time defined in (2.6). Assume $\tau_\infty^\alpha(\omega) > T$, then there is $R_0 > \|x\|_\alpha$ such that $\tau_\infty^{\alpha,R_0}(\omega) > T$, where τ_∞^{α,R_0} is also defined in (2.6). In particular $\|\xi_t(\omega)\|_\alpha \leqslant R_0$ for $t \in [0,T]$. Hence

$$\lambda_n^{\beta-2}\max_{[0,T]}|\xi_n(t,\omega)| \leqslant \lambda_n^{\beta-2-\alpha}\max_{[0,T]}|\lambda_n^\alpha\xi_n(t,\omega)| \leqslant \\ \leqslant \lambda_n^{\beta-2-\alpha}\sup_{[0,T]}\|\xi_t(\omega)\|_\alpha \leqslant R_0\lambda_n^{\beta-2-\alpha}$$

and $\omega \in K_T$.

Vice versa, assume that $\omega \in K_T$ and choose a decreasing sequence $(M_n)_{n\geqslant 1}$ such that $M_n \downarrow 0$ and $\lambda_n^{\beta-2}\max_{[0,T]}|\xi_n(t)| \leqslant M_n$. Set $u_n = \lambda_n^\alpha\xi_n$ and $m_n = \max_{[0,T]}|u_n(t)|$, then

$$\begin{aligned}|u_n(t)| &\leqslant |u_n(0)| + \lambda_n^\alpha|Z_n(t)| + M_{n-1}\lambda_n^2\int_0^t e^{-\nu\lambda_n^2(t-s)}\big(\lambda^{\alpha-2}|u_{n-1}| + \lambda^{2-\beta}|u_n|\big)\,ds\\ &\leqslant |u_n(0)| + \big(\sup_{[0,T]}\lambda_n^\alpha|Z_n(t)|\big) + \frac{1}{\nu}\lambda^{\alpha-2}M_{n-1}m_{n-1} + \frac{1}{\nu}\lambda^{2-\beta}M_{n-1}m_n\end{aligned}$$

and in conclusion

$$\Big(1 - \frac{1}{\nu}\lambda^{2-\beta}M_{n-1}\Big)m_n \leqslant |u_n(0)| + \big(\sup_{[0,T]}\lambda_n^\alpha|Z_n(t)|\big) + \frac{1}{\nu}\lambda^{\alpha-2}M_{n-1}m_{n-1}.$$

For n large enough we have that

$$M_{n-1} \leqslant \frac{\nu}{2\lambda^{2-\beta} + 2\lambda^{\alpha-2}} \qquad\text{and}\qquad \frac{\lambda^{\alpha-2}M_{n-1}}{\nu - \lambda^{2-\beta}M_{n-1}} \leqslant \frac{1}{2},$$

and in particular $\big(1 - \frac{1}{\nu}\lambda^{2-\beta}M_{n-1}\big) \geqslant \frac{1}{2}$, hence

$$m_n \leqslant A_n + \frac{1}{2}m_{n-1}$$

where $A_n = 2\lambda_n^\alpha|x_n| + 2\big(\sup_{[0,T]}\lambda_n^\alpha|Z_n(t)|\big)$. By Lemma 2.6 (applied to an $\alpha' > \alpha$) we have $\sum_n A_n^2 < \infty$ with probability 1, since $\sum A_n^2 \leqslant c(\|x\|_\alpha^2 + \sup_{[0,T]}\|Z\|_{\alpha'}^2)$. Moreover

$$m_n^2 \leqslant A_n^2 + \frac{1}{4}m_{n-1}^2 + A_n m_{n-1} \leqslant \frac{3}{2}A_n^2 + \frac{3}{4}m_{n-1}^2,$$

and by solving the recursive inequality we get $\sum_n m_n^2 < \infty$, which in particular implies that $\tau_\infty^\alpha(\omega) > T$. □

In order to prove Theorem 4.2, we essentially prove that, given a smooth x and $T > 0$, there is a solution $\mathbb{P}_x$ starting at x that satisfies $\mathbb{P}_x[K_T] = 1$, and hence is the unique solution.

Proof of Theorem 4.2. Fix $\alpha \in (\beta-2, \alpha_0+1)$, $x \in V_\alpha$, $T>0$ and an energy martingale solution $\mathbb{P}_x$ starting at x, and let τ_∞^α be the random time defined in (2.6). Without loss of generality we can assume that $\mathbb{P}_x$ is a Galerkin solution (see Definition 2.10). Indeed, by Theorem 2.9 the random time τ_∞^α coincides a. s. with the lifespan τ_x^α of the strong solution with the same initial condition x, and the probability of the event $\{\tau_\infty^x\}$ does not depend on $\mathbb{P}_x$, but only on the strong solution.

Since $\mathbb{P}_x$ is a Galerkin solution, there are $(x^{(N_k)}, \mathbb{P}^{(N_k)})_{k\in\mathbf{N}}$ such that $x^{(N_k)} \to x$ in H and $\mathbb{P}^{(N_k)} \rightharpoonup \mathbb{P}_x$ in Ω_β, and for each k, $\mathbb{P}^{(N_k)}$ is the solution of (2.7), with dimension N_k and initial condition $x^{(N_k)}$. By definition we also have that $x_n^{(N_k)} = x_n$ for $n \leqslant N_k$.

By a standard argument (Skorokhod's theorem) there are a common probability space $(\widetilde{\Omega}, \widetilde{\mathscr{F}}, \widetilde{\mathbb{P}})$ and random variables $X^{(N_k)}$, X on $\widetilde{\Omega}$ with distributions $\mathbb{P}^{(N_k)}$, $\mathbb{P}_x$ respectively, such that, in particular, $X_n^{(N_k)} \to X_n$, $\widetilde{\mathbb{P}}$–a. s., uniformly on $[0,T]$ for all $n \geqslant 1$ and all $T>0$.

Let $\epsilon>0$ be small enough such that $\alpha > \beta-2+2\epsilon$ and $6-2\beta-3\epsilon>0$. We will use Lemma 3.2 with $a_0 < \frac{1}{2}$ (to be chosen later in the proof) and $c_0 = \frac{1}{3}a_0$. Let $\overline{n}$ be the smallest integer such that $\lambda_{n-1}^\alpha |x_n| \leqslant a_0\nu$ for all $n \geqslant \overline{n}$, and set $N_0 = \overline{n} \vee N_{\beta-2+2\epsilon, c_0}(T)$.

For each integer $n_0 \geqslant 1$ and real $M>0$ define the sets

$$A_M(n_0) = \Big\{\sup_{[0,T]}\big(|X_{n_0-1}^{(N_k)}| + |X_{n_0}^{(N_k)}|\big) \leqslant M \text{ for all } k \text{ such that } N_k \geqslant n_0\Big\},$$

Notice that $\widetilde{\mathbb{P}}\big[\bigcup_{M>0} A_M(n_0)\big] = 1$ since $X_n^{(N_k)} \to X_n$ uniformly, $\widetilde{\mathbb{P}}$–a. s., for all $n \geqslant 1$, hence

$$\widetilde{\mathbb{P}}\Big[\bigcup_{n_0\geqslant 1,\ M>0} \big(\{N_0 = n_0\} \cap A_M(n_0)\big)\Big] = 1.$$

Fix $n_0 \geqslant 1$, $M>0$ and $N \geqslant n_0$, then everything boils down to prove that K_T happens on $\{N_0 = n_0\} \cap A_M(n_0)$ for $(X_n^{(N_k)})_{n_0 \leqslant n \leqslant N_k}$ uniformly in k. In the following we work path–wise for $\omega \in \{N_0 = n_0\} \cap A_M(n_0)$ and we aim to use the method in [7], which is based on proving that the area shown in Figure 1 is invariant for the dynamics of the (suitably rescaled) solution. In [7] the area A in the picture is defined for $\beta \leqslant \frac{5}{2}$ with the values $c = \lambda^{-(6-2\beta-3\epsilon)}$, $g(x) = \min(mx+\theta, 1)$, $h(x) = c(\max(x-\delta,0)/(1-\delta))^{\lambda^2}$, $\delta = \frac{1}{10}$, $\theta = \frac{3}{5}$ and $m = \frac{3}{4}$. To deal with the random perturbation we shall need to slightly modify the set A, in particular we will replace the border delimited by $h(x)$ with a new function h_η (see (4.6)).

First we define the re–scaling. Let $\epsilon_n = \nu\lambda_{n-1}^{-2\epsilon}$ and

$$\begin{cases} u_n = \lambda_n^\epsilon(\lambda_n^{\beta-2}Y_n^{(N_k)} + a_0\epsilon_n), \\ v_n = \lambda_n^\epsilon(a_0\epsilon_n - \lambda_n^{\beta-2}Z_n), \end{cases}$$



FIGURE 1. The invariant area.

then we know by Lemma 3.2 that

$$\begin{cases} u_n \geqslant 0, \\ \frac{2}{3}a_0\lambda_n^\epsilon \epsilon_n \leqslant v_n \leqslant \frac{5}{3}a_0\lambda_n^\epsilon \epsilon_n \end{cases} \qquad \text{for all } n_0 \leqslant n \leqslant N_k. \tag{4.1}$$

We have that

$$\begin{aligned}\dot{u}_n = \lambda_n^{2-\epsilon}\Big[-\nu\lambda_n^\epsilon u_n + a_0\nu\lambda_n^{2\epsilon}\epsilon_n + \lambda^{\beta-4+2\epsilon}(u_{n-1}-v_{n-1})^2 + \\ - \lambda^{2-\beta-\epsilon}(u_n - v_n)(u_{n+1}-v_{n+1})\Big].\end{aligned}$$

Define $\delta_0 > 0$ as

$$\frac{1}{\delta_0} = \max\Big\{\frac{1}{\delta}, \lambda_{n_0}^{\beta-2+\epsilon}M + 2a_0\lambda_{n_0}^\epsilon\epsilon_{n_0-1}, \sup_{n\geqslant n_0}\lambda_n^\epsilon(\lambda_n^{\beta-2}x_n + 2a_0\epsilon_n)\Big\}$$

and replace u_n with $U_n(t) = \delta_0^2 u_n(\delta_0 t)$. It turns out that $U_n(0) \leqslant \delta_0 \leqslant \delta$ for all $n \geqslant n_0$ and $\max_{[0,T]} U_{n_0-1} \leqslant \delta_0 \leqslant \delta$ and $\max_{[0,T]} U_{n_0} \leqslant \delta_0 \leqslant \delta$, since

$$|U_{n_0}| \leqslant \delta_0^2\lambda_n^\epsilon\big(\lambda_{n_0}^{\beta-2}|X_{n_0}^{(N_k)}| + a_0\epsilon_{n_0} + \lambda_{n_0}^{\beta-2}|Z_{n_0}|\big) \leqslant \delta_0^2\big(\lambda_{n_0}^{\beta-2+\epsilon}M + 2a_0\lambda_{n_0}^\epsilon\epsilon_{n_0}\big) \leqslant \delta$$

(and similarly for U_{n_0-1}). Moreover the U_n satisfy

$$\dot{U}_n = \lambda_n^{2-\epsilon}\Big[\delta_0^3 P_n^0(U,V) + \delta_0 P_n^1(U,V) + \frac{1}{\delta_0}\lambda^{\beta-4+2\epsilon}P_n^2(U,V)\Big],$$

where

$$\begin{aligned}
P_n^0 &= a_0\nu\lambda_n^{2\epsilon}\epsilon_n + \lambda^{\beta-4+2\epsilon}V_{n-1}^2 - \lambda^{2-\beta-\epsilon}V_nV_{n+1},\\
P_n^1 &= -\nu\lambda_n^\epsilon U_n - 2\lambda^{\beta-4+2\epsilon}V_{n-1}U_{n-1} + \lambda^{2-\beta-\epsilon}(V_{n+1}U_n + V_nU_{n+1}),\\
P_n^2 &= U_{n-1}^2 - \lambda^{6-2\beta-3\epsilon}U_nU_{n+1},
\end{aligned}$$

and $V_n(t) = v_n(\delta_0 t)$.

We proceed now as in [7] and consider for each $n \geqslant n_0$ the coupled systems in (U_n, U_{n+1}),

$$\frac{d}{dt}\begin{pmatrix} U_n \\ U_{n+1} \end{pmatrix} = \lambda_n^{2-\epsilon}\Big(\delta_0^3\mathfrak{B}_n^0 + \delta_0\mathfrak{B}_n^1 + \frac{1}{\delta_0}\lambda^{\beta-4+2\epsilon}\mathfrak{B}_n^2\Big) \tag{4.2}$$

where

$$\mathfrak{B}_n^i = \begin{pmatrix} P_n^i \\ \lambda^{2-\epsilon}P_{n+1}^i \end{pmatrix}, \qquad i = 0, 1, 2,$$

The goal is to prove that $(U_n(t))_{n\geqslant n_0}$ is uniformly bounded in n and t. Indeed, we will show that $0 \leqslant U_n(t) \leqslant 1$ for all $n \in \{n_0, \dots, N_k\}$, which in turns implies that

$$-\lambda_n^\epsilon \epsilon_n \leqslant \lambda_n^{\beta-2+\epsilon} Y_n^{(N_k)} \leqslant \frac{1}{\delta_0^2}.$$

for all $n \in \{n_0, \dots, N_k\}$ and all $t \in [0, T]$. Since $Y_n^{(N_k)} \to Y_n$ uniformly on $[0, T]$ for each n, the same bound holds for the limit Y and, due to Lemma 2.6, $X \in K_T$.

It remains to show that $U_n \leqslant 1$, and to this end we show that each pair $(U_n(t), U_{n+1}(t))$, for $n \geqslant n_0$, remains in A (as it is already in the interior of A at time $t = 0$ by the choice of δ_0). This will be again a dynamical system proof and it is enough to prove that the vector field in (4.2) which gives $(\dot U_n, \dot U_{n+1})$ points inwards on the boundary of A, or equivalently that the scalar product of the vector fields with the normal inwards is positive.

We make two preliminary remarks: first there is nothing to prove for the two pieces of boundary corresponding to the normal vectors $\vec n_1$ and $\vec n_6$, as this is essentially already ensured by Lemma 3.2, and second that since A is convex it is also possible to check that the product of each of the three vector fields $\mathfrak{B}_n^0$, $\mathfrak{B}_n^1$ and $\mathfrak{B}_n^2$ separately is positive.

The vector field $\mathfrak{B}^1$. We will use (4.1), that $U \leqslant 1$ in A and that ϵ_n is non–increasing. We will obtain lower bounds that are positive if the number a_0 is chosen small enough (with size depending only on the constants m, β and ϵ, but not on M, n_0 or δ_0).

On the border with normal $\vec n_2 = (g', 1) = (m, -1)$ we have $U_n \in [0, \frac{1-\theta}{m}]$ and $U_{n+1} = mU_n + \theta$, hence $\lambda^2 U_{n+1} - mU_n \geqslant \lambda^2\theta$ and

$$\begin{aligned} \mathfrak{B}_n^1 \cdot \vec n_2 &= mP_n^1 - \lambda^{2-\epsilon}P_{n+1}^1 \\ &\geqslant \nu\lambda_n^\epsilon(\lambda^2 U_{n+1} - mU_n) - 4a_0(m\lambda^{\beta-4+\epsilon} + \lambda^{4-\beta})\lambda_n^\epsilon\epsilon_{n-1} \\ &\geqslant \lambda_n^\epsilon\big(\nu\lambda^2\theta - a_0 c(m, \beta, \epsilon)\epsilon_{n-1}\big) \end{aligned} \tag{4.3}$$

On the border with normal $\vec n_3 = (0, -1)$ we have $U_{n+1} = 1$, hence

$$\mathfrak{B}_n^1 \cdot \vec n_3 = -\lambda^{2-\epsilon}P_{n+1}^1 \geqslant \lambda^{2-\epsilon}\lambda_{n+1}^\epsilon(\nu - 4a_0\lambda^{2-\beta}\epsilon_{n+1}) \tag{4.4}$$

Similarly, on the border with normal $\vec n_4 = (-1, 0)$ we have $U_n = 1$, hence

$$\mathfrak{B}_n^1 \cdot \vec n_4 = -P_n^1 \geqslant \lambda_n^\epsilon(\nu - 4a_0\lambda^{2-\beta}\epsilon_n). \tag{4.5}$$

Finally we consider the border with normal $\vec{n}_5$. Prior to this we give the precise definition of this piece of the boundary. For $\eta \in (0,1)$ define $\varphi_\eta(x) = \big((x-\eta)/(1-\eta)\big)^{\lambda^2}$, $x \in [\eta, 1]$, and notice that φ_η is positive, increasing, convex and

$$x\varphi_\eta' - \lambda^2\varphi_\eta \geqslant \frac{\lambda^2\eta}{1-\eta}\Big(\frac{\delta-\eta}{1-\eta}\Big)^{\lambda^2-1} > 0.$$

In [7] it was used $h(x) = c\varphi_\delta(x)$. Here we consider

$$h_\eta(x) = \frac{c}{1-\varphi_\eta(\delta)}\big(\varphi_\eta(x) - \varphi_\eta(\delta)\big), \qquad x \in [\delta, 1], \tag{4.6}$$

with $\eta < \delta$ and we notice that h_η is positive, increasing, convex, $h_\eta(\delta) = 0$, $h_\eta(1) = c$ and $h_\eta \to h$ in $C^1([\delta,1])$ as $\eta \uparrow \delta$. Moreover,

$$xh_\eta' - \lambda^2 h_\eta \geqslant \frac{c\lambda^2\delta}{1-\varphi_\eta(\delta)}\,\frac{(\delta-\eta)^{\lambda^2-1}}{(1-\eta)^{\lambda^2}} = c_{\delta-\eta} > 0.$$

With this inequality in hands, we proceed with the estimate of $\mathfrak{B}^1_n \cdot \vec{n}_5$. On the border with normal $\vec{n}_5 = (-h_\eta'(U_n), 1)$ we have $U_n \in [\delta, 1]$ and $U_{n+1} = h_\eta(U_n)$, hence

$$\begin{aligned} \mathfrak{B}^1_n \cdot \vec{n}_5 &= \lambda^{2-\epsilon}P^1_{n+1} - h_\eta'(U_n)P^1_n \\ &\geqslant \lambda_n^\epsilon\big[\nu\big(h_\eta'(U_n)U_n - \lambda^2 h_\eta(U_n)\big) - 4a_0\lambda^{\beta-2+\epsilon}\epsilon_n - 4a_0\lambda^{2-\beta-\epsilon}h'(U_n)\epsilon_n\big] \\ &\geqslant \lambda_n^\epsilon\big(\nu c_{\delta-\eta} - a_0 c(\beta,\epsilon,\delta)\epsilon_n\big), \end{aligned} \tag{4.7}$$

since $h_\eta' \leqslant c\lambda^2/(1-2\delta)$.

The vector field $\mathfrak{B}^2$. The fact that $\mathfrak{B}^2_n \cdot \vec{n}_3$ and $\mathfrak{B}^2_n \cdot \vec{n}_4$ are positive follows immediately from [7], since in those computations the boundary function h is not involved. As it regards $\mathfrak{B}^2_n \cdot \vec{n}_2$ and $\mathfrak{B}^2_n \cdot \vec{n}_5$, it is easy to see that they are both continuous functions of h and h' and with the values of the parameters chosen in [7] (as well as here), both scalar products have positive minima. Hence the same is true if we replace h with h_η, for η small enough, since $h_\eta \to h$ in $C^1([\delta,1])$.

The vector field $\mathfrak{B}^0$. Using (4.1) we have that

$$\begin{aligned} |P^0_n| &\leqslant a_0\nu\lambda_n^{2\epsilon}\epsilon_n + \lambda^{\beta-4+2\epsilon}V_{n-1}^2 + \lambda^{2-\beta-\epsilon}V_nV_{n+1} \\ &\leqslant \lambda_n^\epsilon\big[a_0\nu\lambda_n^\epsilon\epsilon_n + 4a_0^2(\lambda^{\beta-4} + \lambda^{2-\beta})\lambda_n^\epsilon\epsilon_{n-1}^2\big] \end{aligned}$$

and the quantity above can be made a small fraction of λ_n^ϵ if a_0 is small enough. This ensures that together $\mathfrak{B}^1_n$ and $\mathfrak{B}^0_n$ point inward, since the lower bounds for $\mathfrak{B}^1_n$ in formulae (4.3), (4.4), (4.5), (4.7) are strictly positive and independent of n, for a suitable choice of a_0 (up to the common multiplicative factor λ_n^ϵ).

The proof we have given (due to the choice of the numbers m, θ, δ) works for $\beta \leqslant \frac{5}{2}$, hence we can consider β_c slightly larger than $\frac{5}{2}$. A larger value of β_c may be considered (see [7, Remark 2.2]). □

5. THE BLOW–UP TIME

In this section we analyse with more details the blow–up time introduced in Definition 2.3. We give some general results which hold beyond the dyadic model (1.1) under examination and which will be used in the next section to prove that blow–up happens with probability 1. Example 5.6 shows that the fact that blow–up happens almost surely is a property that in general depends on the structure of the drift and hence strongly motivates our analysis.

Thus, consider the local strong solution $(X(\cdot;x),\tau_x)_{x\in\mathcal{W}}$ of a stochastic equation (in finite or infinite dimension) on the state space $\mathcal{W}$, where $\mathcal{W}$ is a separable Hilbert space. By comparing with our case, we assume that

- $\mathbb{P}[\tau_x > 0] = 1$ for all $x \in \mathcal{W}$,
- $X(\cdot;x)$ is continuous for $t < \tau_x$ with values in $\mathcal{W}$,
- $X(\cdot;x)$ is the maximal local solution, namely either $\tau_x = \infty$ or $\|X(t;x)\|_{\mathcal{W}} \to \infty$ as $t \uparrow \infty$, $\mathbb{P}$–a. s.,
- $(X(\cdot;x),\tau_x)_{x\in\mathcal{W}}$ is Markov (in the sense given in Theorem 2.4),
- all martingale solutions coincide with the strong solution up to the blow–up time τ_x.

The last statement plainly implies that the occurrence of blow–up is an intrinsic property of the local strong solution and does not depend in an essential way from weak solutions.

Define for $x \in \mathcal{W}$ and $t \geqslant 0$,

$$\flat(t,x) = \mathbb{P}[\tau_x > t],$$

then we know that $\flat(0,x) = 1$ and $\flat(\cdot,x)$ is non–increasing. Set

$$\flat(x) = \inf_{t\geqslant 0} \flat(t,x) = \mathbb{P}[\tau_x = \infty].$$

We prove a dichotomy for the supremum of $\flat$, namely this quantity is either equal to 0 or to 1.

Lemma 5.1. *Consider the family of processes* (X,τ) *on* $\mathcal{W}$ *as above. If there is* $x_0 \in \mathcal{W}$ *such that* $\mathbb{P}[\tau_{x_0} = \infty] > 0$, *then*

$$\sup_{x\in\mathcal{W}} \mathbb{P}[\tau_x = \infty] = 1$$

Proof. By the Markov property,

$$\flat(t+s,x) = \mathbb{P}[\tau_x > t+s] = \mathbb{E}[\mathbb{1}_{\{\tau_x > t\}}\flat(s, X(t;x))]$$

and in the limit as $s \uparrow \infty$, by monotone convergence,

$$\flat(x) = \mathbb{E}[\mathbb{1}_{\{\tau_x > t\}}\flat(X(t;x))]. \tag{5.1}$$

Set $c = \sup \flat(x)$, then by the above formula,

$$\flat(x_0) = \mathbb{E}[\mathbb{1}_{\{\tau_{x_0} > t\}}\flat(X(t;x_0))] \leqslant c\mathbb{E}[\mathbb{1}_{\{\tau_{x_0} > t\}}] = c\flat(t,x_0),$$

and, as $t \uparrow \infty$, we get $\flat(x_0) \leqslant c\flat(x_0)$, that is $c \geqslant 1$, hence $c = 1$. □

Remark 5.2. If we have the additional information that there is $x_0 \in \mathcal{W}$ such that $\flat(x_0) = 1$, then something more can be said. Indeed, $\mathbb{1}_{\{\tau_{x_0} > t\}} = 1$ almost surely, and, using again formula (5.1),

$$\mathbb{E}[\flat(X(t; x_0))] = \mathbb{E}[\mathbb{1}_{\{\tau_{x_0} > t\}} \flat(X(t; x_0))] = \flat(x_0) = 1,$$

hence $\flat(X(t; x_0)) = 1$, almost surely for every $t > 0$. This is very close to prove that $\flat \equiv 0$, and in fact [28, Theorem 6.8] proves, although with a completely different approach, that under the assumptions of

- (suitable) strong Feller regularity,
- *conditional irreducibility*, namely for every $x \in \mathcal{W}$, $t > 0$ and every open set A in $\mathcal{W}$, $\mathbb{P}[X(t; x) \in A, \tau_x > t] > 0$,

$\flat(x_0) = 1$ implies that $\flat \equiv 1$ on $\mathcal{W}$.

Proposition 5.3. *Consider the family (X, τ) of processes as above. Assume that, given $x \in \mathcal{W}$, there are a closed set $B_\infty \subset \mathcal{W}$ with non–empty interior and three numbers $p_0 \in (0, 1)$, $T_0 > 0$ and $T_1 > 0$ such that*

- $\mathbb{P}[\sigma_{B_\infty}^{x,T_1} = \infty, \tau_x = \infty] = 0$,
- $\mathbb{P}[\tau_y \leqslant T_0] \geqslant p_0$ *for every* $y \in B_\infty$,

where the (discrete) hitting time $\sigma_{B_\infty}^{x,T_1}$ of B_∞, starting from x, is defined as

$$\sigma_{B_\infty}^{x,T_1} = \min\{k \geqslant 0 : X(kT_1; x) \in B_\infty\},$$

and $\sigma_{B_\infty}^{x,T_1} = \infty$ if the set is empty. Then

$$\mathbb{P}[\tau_x < \infty] \geqslant \frac{p_0}{1 + p_0}.$$

Remark 5.4. The first condition in the above proposition can be interpreted as a conditional recurrence: knowing that the solution does not explode, it will visit B_∞ in a finite time with probability 1.

Proof. The first assumption says that $\mathbb{P}[\sigma_{B_\infty}^{x,T_1} > n, \tau_x > n] \downarrow 0$ as $n \to \infty$. If $\mathbb{P}[\tau_x = \infty] = 0$, then $\mathbb{P}[\tau_x < \infty] = 1 \geqslant \frac{p_0}{1+p_0}$ and there is nothing to prove. If on the other hand $\mathbb{P}[\tau_x = \infty] > 0$, then $\mathbb{P}[\tau_x > n] > 0$ for all $n \geqslant 1$ and, since $\mathbb{P}[\tau_x > n] \downarrow \mathbb{P}[\tau_x = \infty]$ as $n \to \infty$, we have that

$$\mathbb{P}[\sigma_{B_\infty}^{x,T_1} \leqslant n | \tau_x > n] = 1 - \frac{\mathbb{P}[\sigma_{B_\infty}^{x,T_1} > n, \tau_x > n]}{\mathbb{P}[\tau_x > n]} \longrightarrow 1, \qquad n \to \infty.$$

For $n \geqslant 1$,

$$\mathbb{P}[\tau_x > n + T_0] \leqslant \mathbb{P}[\tau_x > n + T_0,\ \sigma_{B_\infty}^{x,T_1} \leqslant n] + \mathbb{P}[\sigma_{B_\infty}^{x,T_1} > n].$$

The strong solution is Markov, hence

$$\begin{aligned}\mathbb{P}[\tau_x > n + T_0,\ \sigma^{x,T_1}_{B_\infty} \leqslant n] &= \sum_{k=0}^{n} \mathbb{P}[\tau_x > n + T_0,\ \sigma^{x,T_1}_{B_\infty} = k]\\ &\leqslant (1-p_0)\mathbb{P}[\sigma^{x,T_1}_{B_\infty} \leqslant n].\end{aligned}$$

In conclusion

$$\begin{aligned}\mathbb{P}[\tau_x > n + T_0] &\leqslant (1-p_0)\mathbb{P}[\sigma^{x,T_1}_{B_\infty} \leqslant n] + \mathbb{P}[\sigma^{x,T_1}_{B_\infty} > n]\\ &= 1 - p_0\mathbb{P}[\sigma^{x,T_1}_{B_\infty} \leqslant n]\\ &\leqslant 1 - p_0\mathbb{P}[\sigma^{x,T_1}_{B_\infty} \leqslant n | \tau_x > n]\mathbb{P}[\tau_x > n]\end{aligned}$$

and in the limit as $n \to \infty$,

$$\mathbb{P}[\tau_x = \infty] \leqslant 1 - p_0\mathbb{P}[\tau_x = \infty],$$

that is $\mathbb{P}_x[\tau_x = \infty] \leqslant \frac{1}{1+p_0}$. □

Corollary 5.5. *Assume that there are* $p_0 \in (0,1)$, $T_0 > 0$ *and* $B_\infty \subset \mathcal{W}$ *such that the assumptions of the previous proposition hold for every* $x \in \mathcal{W}$ *(the time* T_1 *may depend on* x*). Then for every* $x \in \mathcal{W}$,

$$\mathbb{P}[\tau_x < \infty] = 1.$$

Proof. The previous proposition yields

$$\sup_{x\in\mathcal{W}} \mathbb{P}[\tau_x = \infty] \leqslant \frac{1}{1+p_0} < 1,$$

hence the dichotomy of Lemma 5.1 immediately implies that $\mathbb{P}[\tau_x < \infty] = 1$ for every $x \in \mathcal{W}$. □

Example 5.6. Here we analyse a simple one dimensional example that shows that the fact that blow–up occurs with full probability does depend on the structure of the drift. We remark that the one dimensional case is fully understood (see for instance [32]), our proofs below are elementary and mimic the proofs of the next section.

Consider the one–dimensional SDEs

$$dX = f_i(X)\,dt + dW, \qquad i = 1,2,$$

with initial condition $X(0) = x \in \mathbf{R}$, where

$$f_1(x) = \begin{cases} x^2, & x \geqslant 0,\\ x, & x < 0,\end{cases} \qquad f_2(x) = \begin{cases} x^2, & x \geqslant 0,\\ -x, & x < 0.\end{cases}$$

It is easy to see (using the Feller test, see for instance [32, Proposition 5.22]) that with the drift f_1 the blow–up function $\flat$ defined in the proof of Lemma 5.1 verifies $0 < \flat(x) < 1$, while with the drift f_2 the blow–up function is $\flat(x) \equiv 0$.

In view of the results proved above and the analysis of the next section (see Theorem 6.1), we observe that

- with both drifts f_1, f_2 and with $B_\infty = \{x \geqslant 1\}$, there are $p_0 \in (0,1)$ and $T_0 > 0$ such that $\mathbb{P}[\tau_x \leqslant T_0] \geqslant p_0$ for all $x \in B_\infty$ (in other words, the second assumption of Proposition 5.3 holds),
- the first assumption of Proposition 5.3 holds for f_2 but *not* for f_1,
- in both cases $\mathbb{E}\big[\sup_{[0,T]}(X_n)_-^p\big] < \infty$ for all $T > 0$ and $p \geqslant 1$.

We give an elementary proof of the first statement, which is modeled on the proof which will be used in the next section for the blow–up of the infinite–dimensional problem (1.1). Given an initial condition $x \in [1, \infty)$, we prove that

$$\mathbb{P}\Big[\Big\{\sup_{t\in[0,2]} |W_t| \leqslant \frac{1}{4}\Big\} \cap \{\tau_x > 2\}\Big] = 0.$$

Indeed, set $Y_t = X_t - W_t$, so that $Y_0 = x$ and

$$dY = dX - dW = (Y + W)^2,$$

in particular $Y_t \geqslant 1$. On the event $\{\sup_{t\in[0,2]} |W_t| \leqslant \frac{1}{4}\}$,

$$\dot{Y} \geqslant Y^2 - 2|W|Y \geqslant Y(Y - \frac{1}{2}) \geqslant \frac{1}{2}Y^2,$$

hence by comparison Y_t (and hence X_t) explodes before time $\frac{2}{x} \leqslant 2$.

6. BLOW–UP FOR $\beta > 3$

In this section we first prove that there are sets in the state space which lead to a blow–up with positive probability. The underlying idea is to use again Lemma 3.2 to adapt the estimates of [10], which work only for positive solutions.

In the second part of the section we show that such sets are achieved with full probability when the blow–up time is conditioned to be infinite. The general result of the previous section immediately implies that blow–up occurs with full probability

6.1. Blow–up with positive probability.

Given $\alpha > \beta - 2$, $p \in (0, \beta - 3)$, $a_0 > 0$ and $M_0 > 0$, define the set

$$B_\infty(\alpha, p, a_0, M_0) = \{x \in V_\alpha : \|x\|_p \geqslant M_0 \text{ and } \inf_{n\geqslant 1} \big(\lambda_{n-1}^{\beta-2} x_n\big) \geqslant -\nu a_0\}. \tag{6.1}$$

We will show that for suitable values of a_0 and M_0 there is a positive probability that the process solution of (1.1) with initial condition in the above set blows up in finite time.

Theorem 6.1. *Let $\beta > 3$ and assume (2.1) and (2.2). Given $\alpha \in (\beta - 2, \alpha_0 + 1)$, $p \in (0, \beta - 3)$, and $a_0 \in (0, \frac{1}{4}]$, there exist $p_0 > 0$, $T_0 > 0$ and $M_0 > 0$ such that* *for each* $x \in B_\infty(\alpha, p, a_0, M_0)$ *and for every energy martingale weak solution* $\mathbb{P}_x$ *with initial condition* x*,*

$$\mathbb{P}_x[\tau_\infty^\alpha \leqslant T_0] \geqslant p_0.$$

Proof. Choose $c_0 > 0$ such that $c_0 \leqslant a_0$ and $c_0 \leqslant \sqrt{a_0}(1-\sqrt{a_0})$, and consider the random integer $N_{\alpha,c_0}(T_0)$ defined in (3.1), where the value of T_0 will be specified at the end of the proof. Set

$$p_0 = \mathbb{P}_x[N_{\alpha,c_0}(T_0) = 1],$$

we remark that $p_0 > 0$ by Lemma 3.1 and its value depends only on the distribution of the solution of (2.4), thus is independent from the martingale solution $\mathbb{P}_x$. The theorem will be proved if we show that

$$\mathbb{P}_x[\tau_\infty^\alpha > T_0,\ N_{\alpha,c_0}(T_0) = 1] = 0, \tag{6.2}$$

indeed

$$\mathbb{P}_x[\tau_\infty^\alpha \leqslant T_0] = 1 - \mathbb{P}_x[\tau_\infty^\alpha \leqslant T_0,\ N_{\alpha,c_0}(T_0) > 1] \geqslant 1 - \mathbb{P}_x[N_{\alpha,c_0}(T_0) > 1] = p_0.$$

We proceed with the proof of (6.2) and we work path–wise on the event

$$\Omega(\alpha, T_0) = \{\tau_\infty^\alpha > T_0\} \cap \{N_{\alpha,c_0}(T_0) = 1\}.$$

Let Z be the solution of (2.4) and $Y = X - Z$ the solution of (2.5). On the event $\{\tau_\infty^\alpha > T_0\}$ (2.5) has a unique solution on $[0, T_0]$, and on the event $\{N_{\alpha,c_0}(T_0) = 1\}$ we have that $\lambda_{n-1}^{\beta-2}|Z_n(t)| \leqslant c_0$ for every $t \in [0, T_0]$ and every $n \geqslant 1$. Set

$$\eta_n = X_n - Z_n + a_0\nu\lambda_{n-1}^{2-\beta},$$

then by this position $\eta = (\eta_n)_{n\geqslant 1}$ satisfies the system

$$\begin{cases} \dot\eta_n = -\nu\lambda_n^2\eta_n + a_0\nu^2\lambda^2\lambda_{n-1}^{4-\beta} + \lambda_{n-1}^\beta X_{n-1}^2 - \lambda_n^\beta X_n X_{n+1}, \\ \eta_n(0) = X_n(0) + a_0\nu\lambda_{n-1}^{2-\beta}. \end{cases} \qquad n \geqslant 1$$

Moreover, by Lemma 3.2 (applied with the values of a_0 and c_0 we have fixed), it turns out that $\eta_n(t;\omega) \geqslant 0$ for each $t \in [0, T_0]$, $n \geqslant 1$ and $\omega \in \Omega(\alpha, T_0)$.

Fix a number $b > 0$, which will be specified later in formula (6.4), then

$$\begin{aligned} \frac{d}{dt}(\eta_n^2 + b\eta_n\eta_{n+1}) &= -2\nu\lambda_n^2\eta_n^2 - b\nu(1+\lambda^2)\lambda_n^2\eta_n\eta_{n+1} \\ &\quad + a_0\lambda^2\nu^2(2 + b\lambda^{4-\beta})\lambda_{n-1}^{4-\beta}\eta_n + a_0 b\lambda^2\nu^2\lambda_{n-1}^{4-\beta}\eta_{n+1} \\ &\quad + 2\lambda_{n-1}^\beta X_{n-1}^2\eta_n + b\lambda_{n-1}^\beta X_{n-1}^2\eta_{n+1} + b\lambda_n^\beta X_n^2\eta_n \\ &\quad - 2\lambda_n^\beta X_n X_{n+1}\eta_n - b\lambda_n^\beta X_n X_{n+1}\eta_{n+1} - b\lambda_{n+1}^\beta\eta_n X_{n+1}X_{n+2} \\ &= \boxed{1} + \boxed{2} + \boxed{3} + \boxed{4}. \end{aligned}$$

By the choice of a_0 and c_0, we have that $(a_0 + c_0)^2 \leqslant a_0$, hence

$$\boxed{2} \geqslant \lambda^2\nu^2(a_0 + c_0)^2(2 + b\lambda^{4-\beta})\lambda_{n-1}^{4-\beta}\eta_n + b\lambda^2\nu^2(a_0 + c_0)^2\lambda_{n-1}^{4-\beta}\eta_{n+1}.$$

The third term on the right–hand side of the formula above can be estimated from below using Young's inequality as

$$\begin{aligned}\boxed{3} &\geqslant 2\lambda_{n-1}^{\beta} X_{n-1}^2 \eta_n + b\lambda_n^{\beta} X_n^2 \eta_n \\ &\geqslant \big[(1+\lambda^{-2p})\lambda_{n-1}^{\beta}\eta_{n-1}^2\eta_n - 4\nu^2(a_0+c_0)^2\frac{\lambda^{2\beta+2p-4}}{\lambda^{2p}-1}\lambda_{n-1}^{4-\beta}\eta_n\big] \\ &\quad + b\big[\lambda_n^{\beta}\eta_n^3 - 2\lambda^{\beta-2}\nu(a_0+c_0)\lambda_n^2\eta_n^2\big]\end{aligned}$$

since $X_n = \eta_n - (a_0\nu\lambda_{n-1}^{2-\beta} - Z_n)$ and $0 \leqslant (a_0\nu\lambda_{n-1}^{2-\beta} - Z_n) \leqslant \nu(a_0+c_0)\lambda_{n-1}^{2-\beta}$. Likewise, the fourth term can be estimated from below as

$$\begin{aligned}\boxed{4} &\geqslant -2\lambda_n^{\beta}\eta_n^2\eta_{n+1} - b\lambda_n^{\beta}\eta_n\eta_{n+1}^2 - b\lambda_{n+1}^{\beta}\eta_n\eta_{n+1}\eta_{n+2} \\ &\quad - \nu^2(a_0+c_0)^2(2\lambda^2 + b\lambda^{6-\beta})\lambda_{n-1}^{4-\beta}\eta_n - b\lambda^2\nu^2(a_0+c_0)^2\lambda_{n-1}^{4-\beta}\eta_{n+1},\end{aligned}$$

where we have dropped the positive terms, using again the fact that $(a_0\nu\lambda_{n-1}^{2-\beta} - Z_n) \geqslant 0$. The three above estimates together yield

$$\frac{d}{dt}\big(\eta_n^2 + b\eta_n\eta_{n+1}\big) + 2\nu\lambda_n^2\eta_n^2 + b\nu(1+\lambda^2)\lambda_n^2\eta_n\eta_{n+1} \geqslant A_n + B_n + C_n,$$

where

$$\begin{aligned}A_n &= b\lambda_n^{\beta}\eta_n^3 + (1+\lambda^{-2p})\lambda_{n-1}^{\beta}\eta_{n-1}^2\eta_n - 2\lambda_n^{\beta}\eta_n^2\eta_{n+1} - b\lambda_n^{\beta}\eta_n\eta_{n+1}^2 \\ &\quad - b\lambda_{n+1}^{\beta}\eta_n\eta_{n+1}\eta_{n+2}, \\ B_n &= -2b\lambda^{\beta-2}\nu(a_0+c_0)\lambda_n^2\eta_n^2, \\ C_n &= -4\nu^2(a_0+c_0)^2\frac{\lambda^{2\beta+2p-4}}{\lambda^{2p}-1}\lambda_{n-1}^{4-\beta}\eta_n.\end{aligned}$$

By using Young's inequality in the same way as [10],
(6.3)

$$\begin{aligned}A_n &\geqslant b\lambda_n^{\beta}\eta_n^3 + (1+\lambda^{-2p})\lambda_{n-1}^{\beta}\eta_{n-1}^2\eta_n - 2\lambda_n^{\beta}\eta_n^2\eta_{n+1} - \frac{1}{2}b\lambda_n^{\beta}\big(\eta_n^2\eta_{n+1} + \eta_{n+1}^3\big) \\ &\quad - \frac{1}{4}b\lambda_{n+1}^{\beta}\big(2\eta_n^2\eta_{n+1} + \eta_{n+1}^2\eta_{n+2} + \eta_{n+2}^3\big) \\ &= b\lambda_n^{\beta}\eta_n^3 - \frac{1}{2}b\lambda_n^{\beta}\eta_{n+1}^3 - \frac{1}{4}b\lambda_{n+1}^{\beta}\eta_{n+2}^3 + (1+\lambda^{-2p})\lambda_{n-1}^{\beta}\eta_{n-1}^2\eta_n \\ &\quad - (2 + \frac{1}{2}b + \frac{1}{2}\lambda^{\beta}b)\lambda_n^{\beta}\eta_n^2\eta_{n+1} - \frac{1}{4}b\lambda_{n+1}^{\beta}\eta_{n+1}^2\eta_{n+2}\end{aligned}$$

Summing up the above estimates yields

$$\begin{aligned}\frac{d}{dt}\Big[\sum_{n=1}^{\infty}\lambda_n^{2p}\left(\eta_n^2+b\eta_n\eta_{n+1}\right)\Big]&+2\nu\|\eta\|_{1+p}^2+\\&+b\nu(1+\lambda^2)\sum_{n=1}^{\infty}\lambda_n^{2+2p}\eta_n\eta_{n+1}\geqslant\sum_{n=1}^{\infty}\lambda_n^{2p}(A_n+B_n+C_n).\end{aligned}$$

By using (6.3) and shifting the summation index we have

$$\begin{aligned}\sum_{n=1}^{\infty}\lambda_n^{2p}A_n&\geqslant b\Big(1-\frac{1}{2}\lambda^{-(\beta+2p)}-\frac{1}{4}\lambda^{-(\beta+4p)}\Big)\sum_{n=1}^{\infty}\lambda_n^{\beta+2p}\eta_n^3\\&\qquad+\Big[\lambda^{2p}-1-b\Big(\frac{1}{2}+\frac{1}{2}\lambda^{\beta}+\frac{1}{4}\lambda^{-2p}\Big)\Big]\sum_{n=1}^{\infty}\lambda_n^{\beta+2p}\eta_n^2\eta_{n+1}\\&=k_1\sum_{n=1}^{\infty}\lambda_n^{\beta+2p}\eta_n^3,\end{aligned}$$

where we have chosen b so that

$$\lambda^{2p}-1-b\Big(\frac{1}{2}+\frac{1}{2}\lambda^{\beta}+\frac{1}{4}\lambda^{-2p}\Big)=0. \tag{6.4}$$

The other two terms are simple, indeed

$$\sum_{n=1}^{\infty}\lambda_n^{2p}B_n=-2b\lambda^{\beta-2}\nu(a_0+c_0)\sum_{n=1}^{\infty}\lambda_n^{2+2p}\eta_n^2=-k_2\|\eta\|_{1+p}^2,$$

and, by the Cauchy–Schwarz inequality,

$$\begin{aligned}\sum_{n=1}^{\infty}\lambda_n^{2p}C_n&=-4\nu^2(a_0+c_0)^2\frac{\lambda^{3\beta+2p-8}}{\lambda^{2p}-1}\sum_{n=1}^{\infty}\lambda_n^{3-\beta+p}(\lambda_n^{1+p}\eta_n)\\&\geqslant-4\nu^2(a_0+c_0)^2\frac{\lambda^{3\beta+2p-8}}{\lambda^{2p}-1}\left(\lambda^{2(3-\beta+p)}-1\right)^{-\frac{1}{2}}\|\eta\|_{1+p}\\&=-k_3\|\eta\|_{1+p},\end{aligned}$$

we have used the fact that $p<\beta-3$. Since on the other hand

$$\begin{aligned}2\nu\|\eta\|_{1+p}^2+b\nu(1+\lambda^2)\sum_{n=1}^{\infty}\lambda_n^{2+2p}\eta_n\eta_{n+1}&\leqslant\nu\big(2+b(1+\lambda^2)\lambda^{-1-p}\big)\|\eta\|_{1+p}^2\\&=k_4\|\eta\|_{1+p}^2,\end{aligned}$$

and

$$\|\eta\|_{1+p}^2=\sum_{n=1}^{\infty}\big(\lambda_n^{\frac{1}{3}(\beta+2p)}\eta_n\big)^2\lambda_n^{-\frac{2}{3}(\beta-3-p)}\leqslant\Big(\frac{1}{k_5}\sum_{n=1}^{\infty}\lambda_n^{\beta+2p}\eta_n^3\Big)^{\frac{2}{3}},$$

all the estimates obtained so far together yield

$$\dot{H} + k_4\psi \geqslant k_1k_5\psi^{\frac{3}{2}} - k_2\psi - k_3\sqrt{\psi},$$

where

$$H(t) = \sum_{n=1}^{\infty} \lambda_n^{2p}\big(\eta_n^2 + b\eta_n\eta_{n+1}\big), \qquad \psi(t) = \|\eta\|_{1+p}^2.$$

Finally, $H \leqslant (1 + b\lambda^{-p})\psi = k_6\psi$, and it is easy to show by a simple argument (following for instance the one in [10]) that if

$$H(0) > M_0^2 := \frac{k_6}{k_1k_5}\big(k_4 + k_2 + \sqrt{(k_4+k_2)^2 + 2k_1k_3k_5}\big) \quad\text{and}\quad T_0 > \frac{4k_6^{\frac{3}{2}}}{k_1k_5\sqrt{H(0)}},$$

the solution of the differential inequality given above for H becomes infinite before time T_0. □

6.2. **Ineluctable occurrence of the blow–up.** The theorem in the previous section has shown that if the initial condition is *not too negative* and the noise is *not too strong*, so that the the process is not kicked away from the quasi–positive area, then the deterministic dynamics dominates and the process explodes. In this section we show that the sets that lead to blow–up are conditionally recurrent, in the sense of Proposition 5.3 (see Remark 5.4).

Theorem 6.2. *Let $\beta > 3$ and assume (2.1) and (2.2). Assume moreover that the set $\{n \geqslant 1 : \sigma_n \neq 0\}$ is non–empty. Given $\alpha \in (\beta - 2, 1 + \alpha_0)$, for every $x \in V_\alpha$ and every energy martingale solution $\mathbb{P}_x$ with initial condition x,*

$$\mathbb{P}_x[\tau_\infty^\alpha < \infty] = 1.$$

Before proving the theorem, we need a few preliminary results. Our strategy is to use Corollary 5.5 to prove that blow–up happens with full probability. Hence we need to prove that sets (6.1) satisfy the assumptions of the corollary. The idea is to find conditions on the size of the solution in H and on the structure of the perturbation, so that the system is led to a set (6.1). Lemma 6.4 shows that the negative part of the solution becomes small, while Lemma 6.5 shows that the size of the solution becomes large. Finally, Lemma 6.6 shows that the conditions found happen with full probability, conditional to the absence of blow–up.

We need a slight improvement of Lemma 3.2. That lemma showed that if the noise is *small* from mode $N_{\alpha,c_0}(T)$ on, and if the initial condition is *not too negative* from mode n_0 on, then the system is *not too negative* from mode $\max(n_0, N_{\alpha,c_0}(T))$ on. In the next lemma we show that after a short time, which depends only on the size of the initial condition *in* H, modes becomes *not too negative* and the above result holds from mode $N_{\alpha,c_0}(T)$ on, regardless of the value of n_0.

We prove the result on the event $\{N_{\alpha,c_0}(T)=1\}$, because the control of the H norm, given in the next lemma, is simpler.

Lemma 6.3. *Let* $\beta>3$ *and assume* (2.1) *and* (2.2)*. There exists* $c_{6.3}>0$ *such that for* $\alpha\in(\beta-2,1+\alpha_0)$*, for every* $x\in V_\alpha$*, every energy martingale solution* $\mathbb{P}_x$ *starting at* x*, every* $T>0$ *and every* $c_0>0$ *with* $4c_0(1+\lambda^{\beta-3})\leqslant 1$*,*

$$\sup_{[0,T]}\|X(t)\|_H\leqslant\|x\|_H+c_{6.3}\nu,$$

$\mathbb{P}_x$*–a. s. on the event* $\{\tau_\infty^\alpha>T\}\cap\{N_{\beta-2,c_0}(T)=1\}$*.*

Proof. As already remarked, since we work on $\{\tau_\infty^\alpha>T\}$, problem (2.5) has a unique solution, so we can work directly on Y (the rigorous proof proceeds through Galerkin approximations). We know that $\lambda_n^{\beta-2}|Z_n(t)|\leqslant c_0\nu$ for $t\in[0,T]$ and $n\geqslant 1$, hence

$$\begin{aligned}\frac{d}{dt}\|Y\|_H^2+2\nu\|Y\|_1^2&\leqslant 2\sum_{n=1}^\infty\lambda_n^\beta(Y_nY_{n+1}Z_n+Y_{n+1}Z_n^2-Y_n^2Z_{n+1}-Y_nZ_nZ_{n+1})\\&\leqslant 2c_0\nu(1+\lambda^{\beta-3})\|Y\|_1^2+2c_0^2\nu^2\frac{\lambda^{\beta-2}}{\lambda^{\beta-3}-1}\|Y\|_1\\&\leqslant 4c_0\nu(1+\lambda^{\beta-3})\|Y\|_1^2+\frac{\lambda^{2\beta-4}}{2(\lambda^{\beta-3}-1)}c_0^3\nu^3.\end{aligned}$$

Using the assumption on c_0 and the fact that $\|Y\|_1\geqslant\lambda\|Y\|_H$, we finally obtain

$$\frac{d}{dt}\|Y\|_H^2+\nu\lambda^2\|Y\|_H^2\leqslant k_0c_0^3\nu^3,$$

where the value of k_0 follows from the inequality above and depends only on β. By integrating the differential inequality, we get

$$\|Y(t)\|_H^2\leqslant\|x\|_H^2+\frac{k_0}{\lambda^2}c_0^3\nu^2,\qquad t\in[0,T],$$

and in conclusion for $t\in[0,T]$,

$$\|X(t)\|_H\leqslant\|Y(t)\|_H+\|Z(t)\|_H\leqslant\|x\|_H+\big(\sqrt{k_0}\lambda^{-1}+(1-\lambda^{4-2\beta})^{-\frac{1}{2}}\big)\nu,$$

where we have used the fact that $c_0\leqslant 1$. □

Lemma 6.4 (Contraction of the negative components)**.** *Let* $\beta>3$ *and assume* (2.1) *and* (2.2)*. For every* $M>0$*,* $a_0\in(0,\frac{1}{4}]$ *and* $c_0<a_0$*, with* $4c_0(1+\lambda^{\beta-3})\leqslant 1$*, there exists* $T_M>0$ *such that for every* $x\in V_\alpha$*, with* $\alpha\in(\beta-2,1+\alpha_0)$ *and* $\|x\|_H\leqslant M$*, and every energy martingale solution* $\mathbb{P}_x$*,*

$$\inf_{n\geqslant 1}\big(\lambda_{n-1}^{\beta-2}X_n(T_M)\big)\geqslant-(a_0+c_0)\nu,$$

$\mathbb{P}_x$*–a. s. on the event* $\{\tau_\infty^\alpha>T_M\}\cap\{N_{\beta-2,c_0}(T_M)=1\}$*.*

Proof. Let n_0 be the first integer such that $\inf_{n\geqslant n_0}\lambda_n^{\beta-2}x_n \geqslant -a_0\nu$. By the choice of c_0 and a_0, we know that $c_0 < \sqrt{a_0}(1-\sqrt{a_0})$, hence Lemma 3.2 implies that $\lambda_n^{\beta-2}Y_n(t) \geqslant -a_0\nu$ holds for every $t\in[0,T_M]$ and every $n\geqslant n_0$. If $n_0 = 1$ there is nothing to prove, so we consider the case $n_0 > 1$.

The idea here is to show that the negative part of the mode n_0-1 becomes closer to 0 within a time T_{n_0-1}. After this time, again by Lemma 3.2, the negative part of the mode n_0-1 stays small, and we can apply the same contraction idea on the negative part of the component n_0-2, which then will become small as well within a time T_{n_0-2}, and so on. The sequence of time intervals turns out to be summable and dependent only on the size of the initial condition in H.

By these considerations it is sufficient to prove the following statement:

> given $n>1$, if we know that for a time $t_0>0$,
>
> $$\sup_{k\geqslant 1}\sup_{[t_0,T]}\lambda_{k-1}^{\beta-2}|Z_k| \leqslant c_0\nu \quad\text{and}\quad \sup_{[t_0,T]}\lambda_n^{\beta-2}(Y_{n+1})_- \leqslant a_0\nu$$
>
> then at time t_0+T_n, where
>
> $$T_n(\|x\|_H, c_0, a_0) = \frac{2}{\nu}(\beta-2)\log\lambda\frac{n-1}{\lambda_n^2} + \frac{2}{\nu}\lambda_n^{-2}\log\Big(1\vee\frac{\|x\|_H + c_{6.3}\nu}{(a_0-c_0)\nu}\Big),$$
>
> we have that
>
> $$Y_n(t_0+T_n) \geqslant -a_0\nu\lambda_{n-1}^{2-\beta}.$$

Before proving the claim, we notice that $\sum_n T_n < \infty$, hence we can choose

$$T_M = \sum_{n=1}^{\infty} T_n(M, c_0, a_0).$$

We conclude the proof of the lemma by showing the above claim. Set

$$\eta_n = Y_n + c_0\nu\lambda_{n-1}^{2-\beta},$$

then $X_n = \eta_n - (c_0\nu\lambda_{n-1}^{2-\beta} - Z_n)$ and

$$\begin{aligned}
\dot\eta_n &= -\nu\lambda_n^2\eta_n + c_0\nu^2\lambda^2\lambda_{n-1}^{4-\beta} + \lambda_{n-1}^{\beta}X_{n-1}^2 - \lambda_n^{\beta}X_nX_{n+1}\\
&\geqslant -(\nu\lambda_n^2 + \lambda_n^{\beta}X_{n+1})\eta_n + c_0\nu^2\lambda^2\lambda_{n-1}^{4-\beta} + \lambda_n^{\beta}(c_0\nu\lambda_{n-1}^{2-\beta} - Z_n)X_{n+1}\\
&\geqslant -(\nu\lambda_n^2 + \lambda_n^{\beta}X_{n+1})\eta_n + c_0\nu^2\lambda^2\lambda_{n-1}^{4-\beta} - \lambda_n^{\beta}(c_0\nu\lambda_{n-1}^{2-\beta} - Z_n)(X_{n+1})_-.
\end{aligned}$$

By the assumption of the claim, $(X_{n+1})_- \leqslant (a_0+c_0)\nu\lambda_n^{2-\beta}$ and $(c_0\nu\lambda_{n-1}^{2-\beta} - Z_n) \leqslant 2c_0\nu\lambda_{n-1}^{2-\beta}$, hence

$$\begin{aligned}
\dot\eta_n &\geqslant -(\nu\lambda_n^2 + \lambda_n^{\beta}X_{n+1})\eta_n + c_0\lambda^2\nu^2\big(1-2(a_0+c_0)\big)\lambda_{n-1}^{4-\beta}\\
&\geqslant -(\nu\lambda_n^2 + \lambda_n^{\beta}X_{n+1})\eta_n,
\end{aligned}$$

which implies that for $t \geqslant t_0$,

$$\begin{aligned}\eta_n(t) &\geqslant \eta_n(t_0)\exp\Big(-\int_{t_0}^t (\nu\lambda_n^2 + \lambda_n^\beta X_{n+1})\,ds\Big) \geqslant\\ &\geqslant -(\eta_n(t_0))_-\exp\Big(-\int_{t_0}^t (\nu\lambda_n^2 + \lambda_n^\beta X_{n+1})\,ds\Big) \geqslant -(\eta_n(t_0))_-\,e^{-\frac{1}{2}\nu\lambda_n^2(t-t_0)},\end{aligned}$$

since, using the fact that $a_0 + c_0 \leqslant \frac{1}{2}$,

$$\nu\lambda_n^2 + \lambda_n^\beta X_{n+1} \geqslant \nu\lambda_n^2 - \nu(a_0+c_0)\lambda_n^2 = \nu\lambda_n^2\big(1-(a_0+c_0)\big) \geqslant \frac{1}{2}\nu\lambda_n^2.$$

Finally, by Lemma 6.3,

$$(\eta_n(t_0))_- \leqslant (Y_n(t_0))_- \leqslant |Y_n(t_0)| \leqslant \|Y(t_0)\|_H \leqslant \|x\|_H + c_{6.3}\nu,$$

and it is elementary now to check that at time $t_0 + T_n$,

$$\begin{aligned}Y_n(t_0+T_n) &= \eta_n(t_0+T_n) - c_0\nu\lambda_{n-1}^{2-\beta} \geqslant\\ &\geqslant -(\|x\|_H + c_{6.3})\,e^{-\frac{1}{2}\nu\lambda_n^2 T_n} - c_0\nu\lambda_{n-1}^{2-\beta} \geqslant -a_0\nu\lambda_n^{2-\beta},\end{aligned}$$

which concludes the proof of the claim. □

In order to show that the hitting time for sets like (6.1), where there is a uniform estimate on the probability of blow–up, is finite, we also need to ensure that the size of the solution is large enough, while being *not too negative*.

At this stage the noise is crucial, although one randomly perturbed component is enough for our purposes. The underlying ideas of the following lemma come from control theory, although we do not need sophisticated results like [45, 39], because a quick and strong impulse turns out to be sufficient.

Lemma 6.5 (Expansion in H). *Under the assumptions of Theorem 6.2, let m be the smallest element of the set $\{n \geqslant 1 : \sigma_n \neq 0\}$. Given $M_1 > 0$, $M_2 > 0$, and $a_0, a_0', c_0 > 0$ such that $c_0 < a_0 < a_0' < \frac{1}{4}$ and $c_0 + a_0 < a_0'$, for every $X(0) \in V_\alpha$, with $\|X(0)\|_H \leqslant M_1$ and $\inf_{n\geqslant 1}\lambda_{n-1}^{\beta-2}X_n(0) \geqslant -a_0\nu$, there exists $T > 0$, whose value depends on M_1, M_2, c_0, a_0, a_0' and m, such that*

- *$\lambda_{n-1}^{\beta-2}X_n(t) \geqslant -(a_0'+c_0)\nu$ for every $n \geqslant 1$ and $t \in [0,T]$,*
- *$\|X(T)\|_H \geqslant M_2$,*

on the event

$$\{\tau_\infty^\alpha > T_M\}\cap\{\sup_{[0,T]}\lambda_{n-1}^{\beta-2}|Z_n(t)| \leqslant c_0\nu \text{ for } n\neq m\}\cap\{\sup_{[0,T]}\lambda_{m-1}^{\beta-2}|Z_m(t)-\psi(t)| \leqslant c_0\nu\},$$

where $\psi : [0,T] \to \mathbf{R}$ is an non–decreasing continuous function such that $\psi(0) = 0$ and $\psi(T)$ large enough depending on the above given data (its value is given in the proof).

Proof. We work on the event given in the statement of the proof.
Step 1: estimate in H. We first give an estimate of X in H. Set $\widetilde{\psi} = \sup_{[0,T]} \|Z_m\|_H \leqslant \psi(T) + c_0\nu$, then

$$\begin{aligned}
\frac{d}{dt}\|Y\|_H^2 + 2\nu\|Y\|_1^2 &\leqslant 2\sum_{n=1}^{\infty} \lambda_n^\beta X_n(Z_n Y_{n+1} - Y_n Z_{n+1}) \\
&\leqslant 2\sum_{n=1}^{\infty} \lambda_n^\beta\big(|Z_n Y_n Y_{n+1}| + Z_n^2|Y_{n+1}| + |Z_{n+1}|Y_n^2 + |Z_n Z_{n+1} Y_n|\big) \\
&\quad + 2\lambda_{m-1}^\beta\big(|Z_m|Y_{m-1}^2 + |Z_{m-1} Z_m Y_{m-1}|\big) \\
&\quad + 2\lambda_m^\beta\big(||Z_m Y_m Y_{m+1} + Z_m^2|Y_{m+1}| + |Z_m Z_{m+1} Y_m|\big) \\
&\leqslant 4c_0\nu(1+\lambda^{\beta-3})\|Y\|_1^2 + k_0 c_0^3 \nu^3 \\
&\quad + \lambda_m^\beta(2\widetilde{\psi}^2 + 4\widetilde{\psi} + 4c_0\nu\lambda^{\beta-3}\widetilde{\psi})(1 + \|Y\|_H^2) \\
&\leqslant \nu\|Y\|_1^2 + k_0\nu^3 + 16\lambda_m^\beta(1+\nu)(1+\widetilde{\psi}^2)(1+\|Y\|_H^2),
\end{aligned}$$

where we have estimated the terms without Z_m as in Lemma 6.3 (the constant k_0 is the same of that lemma) and the terms with Z_m bounding the components of Y with $\|Y\|_H$. If $k_1 = k_0\nu^3$, $k_2 = 16\lambda_m^\beta(1+\nu)$ and $M_3(T,\psi(T))^2 = (M_1^2 + k_1/k_2)\exp\big(k_2 T(1+\widetilde{\psi}^2)\big)$, it follows from Gronwall's lemma that $\sup_{[0,T]}\|Y(t)\|_H^2 \leqslant M_3(T,\psi(T))^2$. Since on the given event we have that $\|Z(t)\|_H \leqslant \lambda^{2-\beta}(\lambda^{2-\beta}-1)^{-1} + c_0\nu + \psi(T)$ for every $t \in [0,T]$, we finally have that

$$\begin{aligned}
\sup_{[0,T]}\|X(t)\|_H &\leqslant c_0\nu + \frac{\lambda^{2-\beta}}{\lambda^{2-\beta}-1} + \psi(T) + \Big(M_1 + \sqrt{\tfrac{k_1}{k_2}}\Big)\, e^{k_2 T(1+(\psi(T)+c_0\nu)^2)} \\
&=: M_4(T,\psi(T))
\end{aligned}$$

Step 2: Large size at time T. Using the previous estimate we have

$$\begin{aligned}
X_m(t) &= e^{-\nu\lambda_m^2 t} X_m(0) + Z_m(t) + \int_0^t e^{-\nu\lambda_m^2(t-s)}\big(\lambda_{m-1}^\beta X_{m-1}^2 - \lambda_m^\beta X_m X_{m+1}\big)\,ds \\
&\geqslant -a_0\nu\lambda_{m-1}^{2-\beta} + \big(\psi(t) - c_0\nu\lambda_{m-1}^{2-\beta}\big) - \lambda_m^\beta t \sup_{[0,T]}\|X\|_H^2,
\end{aligned}$$

hence for $t = T$ we have

$$X_m(T) \geqslant \psi(T) - \nu - \lambda_m^\beta M_4(T,\psi(T))T.$$

We can choose $\psi(T) = M_2 + 2\nu$ and notice that $M_4(T, M_2 + 2\nu)T \to 0$ as $T \downarrow 0$, hence we can choose T small enough so that $\lambda_m^\beta M_4(T,\psi(T))T \leqslant \nu$, to obtain $X_m(T) \geqslant M_2$ and in conclusion $\|X(T)\|_H \geqslant M_2$.
Step 3: Bound from below for $n = m$. The choice of $\psi(T)$ and the computation in the above step yield

$$\lambda_{m-1}^{\beta-2} X_m(t) \geqslant -(a_0 + c_0)\nu - \lambda^{2-\beta} M_4(T, M_2 + 2\nu)\lambda_m^{2\beta-2} T,$$

since ψ is non–negative. By assumption we have that $a_0 + c_0 < a_0'$, hence, possibly fixing a smaller value of T than the one chosen in the previous step, we can ensure that X_m has $-a_0'\nu\lambda_{m-1}^{2-\beta}$ as lower bound on $[0,T]$.
Step 4: Bound from below for $n \neq m$. If $n > m$, the proof proceeds as in Lemma 3.2, since X_m appears in the system of equations for $(Y_n)_{n>m}$ only through the *positive* term $\lambda_m^\beta X_m^2$ in the equation for the $(m+1)^{\text{th}}$ component (and since $c_0 < \sqrt{a_0'}(1-\sqrt{a_0'})$ by the choice of c_0, a_0').

If $n < m$, the proof follows by finite induction. Indeed, the lower bound is true for $n = m$ by the previous step. Next we prove that if $\lambda_n^{2-\beta}Y_{n+1}(t) \geqslant -a_0'\nu$ for all $t \in [0,T]$, then $\lambda_{n-1}^{2-\beta}Y_n \geqslant -a_0'\nu$. We prove this similarly to the proof of Lemma 6.4. Indeed, set $\eta_n = Y_n + a_0'\nu\lambda_{n-1}^{2-\beta}$, then using only that $\lambda_{n-1}^{\beta-2}|Z_n(t)| \leqslant c_0\nu$ and $(X_{n+1}(t))_- \leqslant (a_0' + c_0)\nu\lambda_n^{2-\beta}$ for $t \in [0,T]$,

$$\begin{aligned}\dot\eta_n &\geqslant -(\nu\lambda_n^2 + \lambda_n^\beta X_{n+1})\eta_n + a_0'\nu^2\lambda^{\beta-2}\lambda_n^{4-\beta} - \lambda_n^\beta(a_0'\nu\lambda_{n-1}^{2-\beta} - Z_n)(X_{n+1})_- \\ &\geqslant -(\nu\lambda_n^2 + \lambda_n^\beta X_{n+1})\eta_n + \nu\lambda^{\beta-2}\big(a_0' - (a_0'+c_0)^2\big)\lambda_n^{4-\beta} \\ &\geqslant -(\nu\lambda_n^2 + \lambda_n^\beta X_{n+1})\eta_n,\end{aligned}$$

since $c_0 < a_0' < \frac{1}{4}$, hence $c_0 < \sqrt{a_0'}(1-\sqrt{a_0'})$. The fact that $\eta_n(0) \geqslant 0$ implies that $\eta_n(t) \geqslant 0$. $\square$

Lemma 6.4 and Lemma 6.5 suggest how to have a *well-behaved* noise, namely a random perturbation that leads the system from a ball in H to a pre–blow–up set (6.1). Let $c_0 > 0$, $t_0 > 0$, $T_c > 0$, $T_e > 0$ and $\psi : [0, T_e] \to \mathbf{R}$ be a non–negative non–increasing function, and define the event

$$\mathcal{N}(t_0; c_0, T_c, T_e, \psi) = \mathcal{N}_c(c_0, t_0, T_c) \cap \mathcal{N}_e(c_0, t_0 + T_c, T_e, \psi)$$

where

$$\mathcal{N}_c(c_0, t_0, T_c) = \big\{\lambda_{n-1}^{\beta-2}|Z_n^c(t)| \leqslant c_0\nu \text{ for all } n \geqslant 1 \text{ and } t \in [t_0, t_0 + T_c]\big\}$$

and

$$\begin{aligned}\mathcal{N}_e(c_0, t_0 + T_c, T_e, \psi) = \Big\{ &\sup_{[t_0+T_c, t_0+T_c+T_e]} \lambda_{m-1}^{\beta-2}|Z_m^e(t) - \psi_{t_0+T_c}(t)| \leqslant c_0\nu, \\ &\lambda_{n-1}^{\beta-2}|Z_n^e(t)| \leqslant c_0\nu \text{ for all } n \neq m \text{ and } t \in [t_0+T_c, t_0+T_c+T_e]\Big\},\end{aligned}$$

$\psi_s : [s, s+T_e] \to \mathbf{R}$ is defined for $s \geqslant 0$ as $\psi_s(t) = \psi(t-s)$, for $t \in [s, s+T_e]$, m is the smallest integer of the set $\{n : \sigma_n \neq 0\}$, and for every $n \geqslant 1$,

$$Z_n^c(t) = \sigma_n \int_{t_0}^t e^{-\nu\lambda_n^2(t-t_0)}\, dW_n, \qquad t \in [t_0, t_0 + T_c],$$

$$Z_n^e(t) = \sigma_n \int_{t_0+T_c}^t e^{-\nu\lambda_n^2(t-t_0-T_c)}\, dW_n, \qquad t \in [t_0 + T_c, t_0 + T_c + T_e].$$

Given a weak martingale solution $\mathbb{P}_x$ with initial condition x, the following facts are straightforward:

- $\mathcal{N}_c(c_0, t_0, T_c)$ and $\mathcal{N}_e(c_0, t_0 + T_c, T_e, \psi)$ are independent,
- they both have positive probability (Lemma 3.1 ensures that $\mathcal{N}_c$ has positive probability and the same methods can be used for $\mathcal{N}_e$),
- the value of their probability is independent of t_0,
- if t_0, T_c, T_e and t_0' are given such that $t_0+T_c+T_e \leqslant t_0'$, then $\mathcal{N}(t_0; c_0, T_c, T_e, \psi)$ and $\mathcal{N}(t_0'; c_0, T_c, T_e, \psi)$ are independent.

The basic idea for the proof of Theorem 6.2 is to show that a pre–blow–up set (6.1) is recurrent for the system, knowing that there is no blow–up. Since a *well-behaved* randomness leads the system from a ball in H to a pre–blow–up set, it is sufficient to show that the fact that the system hits a ball in H, of some radius, and *afterwards* the randomness is *well-behaved* happens with probability one.

Lemma 6.6. *Let* $\beta > 3$, *assume* (2.1) *and* (2.2) *and consider* $\alpha \in (\beta-2, \alpha_0+1)$. *There exists* $c_{6.6} > 0$ *such that if* $M > 0$, $T_c > 0$, $T_e > 0$, $c_0 > 0$, *and* $\psi : [0, T_e] \to \mathbf{R}$ *is a non–decreasing and non–negative function and*

$$\frac{c_{6.6}}{M^2} + e^{-\nu\lambda^2(T_c+T_e)} < 1,$$

then for every $x \in V_\alpha$ *and every energy martingale solution* $\mathbb{P}_x$ *starting at* x,

$$\mathbb{P}_x\Big[\{\tau_\infty^\alpha = \infty\} \cap \bigcap_{k\geqslant 1}\Big(\{\|X(kT)\|_H \leqslant M\} \cap \mathcal{N}(kT; c_0, T_c, T_e, \psi)\Big)^c\Big] = 0,$$

where $T = T_c + T_e$.

Proof. We first obtain a quantitative estimate on the return time in balls of H of the Markov process $X^R(\cdot; x)$, solution of problem (2.3) with initial condition $x \in V_\alpha$, which has been introduced in the proof of Theorem 2.4, and then get the same estimate for the strong solution. This will allow us to prove the lemma.
Step 1. It is straightforward to prove by Itô's formula (applied to $\|X^R(t; x)\|_H^2$) and Gronwall's lemma that

$$\mathbb{E}[\|X^R(t; x)\|_H^2] \leqslant \|x\|_H^2\, e^{-2\nu\lambda^2 t} + c_{6.6}, \tag{6.5}$$

where $c_{6.6} = (2\nu\lambda^2)^{-1}\sum_{n=1}^\infty \sigma_n^2$, and where the last series is convergent due to (2.1) and to the assumption $\alpha_0 > \beta - 3$.
Step 2. We use the previous estimate to show that

$$\mathbb{P}\big[\|X^R(kT; x)\|_H \geqslant M \text{ for } k = 1, \dots, n\big] \leqslant \Big(e^{-\nu\lambda^2 T} + \frac{c_{6.6}}{M^2}\Big)^{n-1}. \tag{6.6}$$

We proceed for instance as in [18, Lemma III.2.4], we detail the proof for the sake of completeness. Define, for k integer, $C_k = \{\|X^R(kT; x)\|_H \geqslant M\}$ and

$B_k = \bigcap_{j=0}^k C_j$, and set $\alpha_k = \mathbb{E}[\mathbb{1}_{B_k}\|X^R(kT;x)\|_H^2]$ and $p_k = \mathbb{P}[B_k]$. By the Markov property, Chebychev's inequality and (6.5),

$$\mathbb{P}[C_{k+1}|\mathscr{F}_{kT}] \leqslant \frac{1}{M^2}\,e^{-\nu\lambda^2 T}\,\|X^R(kT;x)\|_H^2 + \frac{c_{6.6}}{M^2},$$

hence

$$p_{k+1} = \mathbb{E}\big[\mathbb{1}_{B_k}\mathbb{P}[C_{k+1}|\mathscr{F}_{kT}]\big] \leqslant \frac{1}{M^2}\,e^{-\nu\lambda^2 T}\,\alpha_k + \frac{c_{6.6}}{M^2}p_k.$$

On the other hand, by integrating (6.5) on B_k, we get

$$\alpha_{k+1} \leqslant \mathbb{E}[\mathbb{1}_{B_k}\|X^R((k+1)T;x)\|_H^2] \leqslant e^{-\nu\lambda^2 T}\,\alpha_k + c_{6.6}p_k$$

If $(\widetilde{\alpha}_k)_{k\in\mathbf{N}}$ and $(\widetilde{p}_k)_{k\in\mathbf{N}}$ are the solutions to the recurrence system

$$\begin{cases}\widetilde{\alpha}_{k+1} = e^{-\nu\lambda^2 T}\,\widetilde{\alpha}_k + c_{6.6}\widetilde{p}_k,\\ \widetilde{p}_{k+1} = \frac{1}{M^2}\,e^{-\nu\lambda^2 T}\,\widetilde{\alpha}_k + \frac{c_{6.6}}{M^2}\widetilde{p}_k,\end{cases}\qquad k \geqslant 1,$$

with $\widetilde{p}_1 = p_1$ and $\widetilde{\alpha}_1 = \alpha_1$, then $\alpha_k \leqslant \widetilde{\alpha}_k$ and $p_k \leqslant \widetilde{p}_k$ for all $k \geqslant 1$ and (6.6) easily follows, since $\widetilde{\alpha}_k = M^2\widetilde{p}_k$ for $k \geqslant 2$.

Step 3. By the previous step,

$$\begin{aligned}&\mathbb{P}[\|X(kT;x)\|_H \geqslant M, k = 1,\dots,n, \tau_x^\alpha = \infty]\\ &\qquad\leqslant \lim_{R\uparrow\infty}\mathbb{P}[\|X(kT;x)\|_H \geqslant M, k = 1,\dots,n, \tau_x^{\alpha,R} > nT]\\ &\qquad= \lim_{R\uparrow\infty}\mathbb{P}[\|X^R(kT;x)\|_H \geqslant M, k = 1,\dots,n, \tau_x^{\alpha,R} > nT]\\ &\qquad\leqslant \limsup_{R\uparrow\infty}\mathbb{P}[\|X^R(kT;x)\|_H \geqslant M, k = 1,\dots,n]\\ &\qquad\leqslant \Big(e^{-\nu\lambda^2 T} + \frac{c_{6.6}}{M^2}\Big)^{n-1},\end{aligned}$$

since $\tau_x^\alpha = \sup_{R>0}\tau_x^{\alpha,R}$. Hence, if we define the (discrete) hitting time $K_1 = \min\{k \geqslant 0 : \|X(kT;x)\|_H \leqslant M\}$ of the ball $B_M(0)$ in H (and $K_1 = \infty$ if the set is empty), then $K_1 < \infty$ on the event $\{\tau_x^\alpha = \infty\}$. Likewise, define the sequence of (discrete) return times $K_2 = \min\{k > K_1 : \|X(kT;x)\|_H \leqslant M\}$, …, $K_j = \min\{k > K_{j-1} : \|X(kT;x)\|_H \leqslant M\}$, …, with the understanding that a return time is infinite if the corresponding set is empty (this in particular happens if some previous return time is already infinite). The previous step immediately implies that $K_j < \infty$ for each $j \geqslant 1$ on the event $\{\tau_x^\alpha = \infty\}$.

Step 4. Consider for $k \geqslant 1$ the events $\mathcal{N}_k = \mathcal{N}(kT; c_0, T_c, T_e, \psi)$. By the considerations we have stated above, we know that $\mathbb{P}[\mathcal{N}_k]$ is constant in k, and we set $p = \mathbb{P}[\mathcal{N}_k]$. Moreover, by the choice of T, it turns out that $\mathcal{N}_1, \mathcal{N}_2, \dots, \mathcal{N}_k, \dots$ are all independent. Set $\mathcal{N}_\infty = \emptyset$ and define the time

$$L_0 = \min\{j \geqslant 1 : \mathbb{1}_{\mathcal{N}_{K_j}} = 1\},$$

and $L_0=\infty$ if the set is empty. Notice that if L_0 is finite, then $\|X(K_{L_0}T;x)\|_H\leqslant M$ and the random perturbation is *well-behaved* in the time interval $[K_{L_0}T, K_{L_0}T+T_c+T_e]$. Hence the lemma is proved if we show that

$$\mathbb{P}[L_0=\infty,\tau_x^\alpha=\infty]=0. \tag{6.7}$$

Step 5. Given an integer $\ell\geqslant 1$, we have that

$$\begin{aligned}\mathbb{P}[L_0>\ell,\tau_x^\alpha=\infty]&=\mathbb{P}[\mathcal{N}^c_{K_1}\cap\dots\mathcal{N}^c_{K_\ell}\cap\{\tau_x^\alpha=\infty\}]\\&=\sum_{k_1=1}^\infty\cdots\sum_{k_\ell=k_{\ell-1}+1}^\infty\mathbb{P}[S_\ell(k_1,\dots,k_l)\cap\{\tau_x^\alpha=\infty\}],\end{aligned}$$

where we have set

$$S_\ell(k_1,\dots,k_\ell)=\mathcal{N}^c_{K_1}\cap\dots\mathcal{N}^c_{K_\ell}\cap\{K_1=k_1,\dots,K_\ell=k_\ell\}.$$

Notice that $S_\ell(k_1,\dots,k_\ell)\in\mathscr{F}_{(k_\ell+1)T}$, hence by the Markov property,

$$\begin{aligned}&\mathbb{P}[S_\ell(k_1,\dots,k_l)\cap\{\tau_x^\alpha>(k_\ell+1)T\}]\\&\quad=\mathbb{E}\Big[\mathbb{1}_{S_{\ell-1}(k_1,\dots,k_{\ell-1})}\mathbb{1}_{\{\tau_x^\alpha>(k_{\ell-1}+1)T\}}\mathbb{1}_{\{K_\ell=k_\ell\}}\mathbb{P}[\mathcal{N}^c_{k_\ell}\cap\{\tau^\alpha_{X(k_\ell T;x)}>T\}|\mathscr{F}_{k_\ell T}]\Big]\\&\quad\leqslant(1-p)\mathbb{P}[S_{\ell-1}(k_1,\dots,k_{\ell-1})\cap\{\tau_x^\alpha>(k_{\ell-1}+1)T\}\cap\{K_\ell=k_\ell\}].\end{aligned}$$

By summing up over k_ℓ, we have

$$\begin{aligned}&\sum_{k_\ell=k_{\ell-1}+1}^\infty\mathbb{P}[S_\ell(k_1,\dots,k_\ell)\cap\{\tau_x^\alpha>(k_\ell+1)T\}]\leqslant\\&\qquad\qquad\leqslant(1-p)\mathbb{P}[S_{\ell-1}(k_1,\dots,k_{\ell-1})\cap\{\tau_x^\alpha>(k_{\ell-1}+1)T\}]\end{aligned}$$

and the iteration of this argument finally yields

$$\mathbb{P}[L_0>\ell,\tau_x^\alpha=\infty]\leqslant(1-p)^\ell,$$

hence the claim (6.7). ☐

We have all the ingredients to complete the proof of the main theorem of this section.

Proof of Theorem 6.2. Fix $\alpha\in(\beta-2,1+\alpha_0)$, $\overline{p}\in(0,\beta-3)$ and $\overline{a_0}\in(0,\frac14]$, and consider the values $\overline{p_0}>0$, and $\overline{M_0}>0$ given by Theorem 6.1. In view of Corollary 5.5, it is enough to prove that the (sampled) arrival time to $B_\infty(\alpha,\overline{p},\overline{a_0},\overline{M_0})$ is finite on the event $\{\tau_x^\alpha=\infty\}$, for all $x\in V_\alpha$. To prove this, by virtue of Lemma 6.6, it is sufficient to prove that there are $M, T_c, T_e, c_0>0$ and a function ψ as in the statement of the lemma such that $e^{-\nu\lambda^2T_c}+\frac{c_{6.6}}{M^2}<1$ and

$$\left.\begin{aligned}\|X(t_0;x)\|_H\leqslant M\\ \mathcal{N}(t_0;c_0,T_c,T_e,\psi)\end{aligned}\right\}\quad\Rightarrow\quad X(t_0+T_c+T_e;x)\in B_\infty(\alpha,\overline{p},\overline{a_0},\overline{M_0}). \tag{6.8}$$

Indeed, the left–hand side of the above implication happens almost surely on $\{\tau_x^\alpha < \infty\}$ for some integer k such that $t_0 = k(T_c + T_e)$, hence the right–hand side happens with the same probability and $\mathbb{P}[\sigma_{B_\infty}^{x,T_c+T_e} = \infty, \tau_x^\infty = \infty] = 0$.

We conclude with the proof of (6.8). We first notice that in Lemma 6.4, the larger we choose M, the larger is the time T_c we get, hence we apply Lemma 6.4 with $a_0 = \frac{1}{8}\overline{a_0}$, $c_0 < \min\{\frac{1}{8}\overline{a_0}, (4(1+\lambda^{\beta-3}))^{-1}\}$ and $M > 0$ large enough so that the time T_c whose existence is ensured by the lemma satisfies $e^{-\nu\lambda^2 T_c} + \frac{c_{6.6}}{M^2} < 1$. Moreover we know that

- $\inf_{n\geqslant 1}\big(\lambda_{n-1}^{\beta-2}X_n(t_0+T_c)\big) \geqslant -(a_0+c_0)\nu \geqslant -\frac{1}{4}\overline{a_0}\nu$ on the event $\{\tau_x^\alpha = \infty\} \cap \mathcal{N}_c(c_0, t_0, T_c)$,
- $\|X(t_0+T_c)\|_H \leqslant \|X(t_0)\|_H + c_{6.3}\nu \leqslant M + c_{6.3}\nu$,

where the second statement follows from Lemma 6.3. We apply then Lemma 6.5 with $M_1 = M + c_{6.3}\nu$, $M_2 = \overline{M_0}$, $a_0 = \frac{1}{4}\overline{a_0}$, $a_0' = 2a_0$ and c_0 as above, hence there is $T_e > 0$ such that

- $\inf_{n\geqslant 1}\big(\lambda_{n-1}^{\beta-2}X_n(t_0+T_c+T_e)\big) \geqslant -(a_0'+c_0)\nu \geqslant -\overline{a_0}\nu$ on the event $\{\tau_x^\alpha = \infty\} \cap \mathcal{N}_e(c_0, t_0+T_c, T_e, \psi)$,
- $\|X(t_0+T_c+T_e)\|_{\overline{p}} \geqslant \lambda^{\overline{p}}\|X(t_0+T_c+T_e)\|_H \geqslant \overline{M_0}$,

that is $X(t_0+T_c+T_e) \in B_\infty(\alpha, \overline{p}, \overline{a_0}, \overline{M_0})$. □

REFERENCES

1. Sergio Albeverio, Arnaud Debussche, and Lihu Xu, *Exponential mixing of the 3D stochastic Navier-Stokes equations driven by mildly degenerate noises*, 2009.
2. David Barbato, Michele Barsanti, Hakima Bessaih, and Franco Flandoli, *Some rigorous results on a stochastic GOY model*, J. Stat. Phys. **125** (2006), no. 3, 673–712.
3. David Barbato, Franco Flandoli, and Francesco Morandin, *Anomalous dissipation in a stochastic inviscid dyadic model*, 2010, arXiv:1007.1004, to appear on Ann. Appl. Probab.
4. ______, *A theorem of uniqueness for an inviscid dyadic model*, C. R. Math. Acad. Sci. Paris **348** (2010), no. 9-10, 525–528.
5. ______, *Uniqueness for a stochastic inviscid dyadic model*, Proc. Amer. Math. Soc. **138** (2010), no. 7, 2607–2617.
6. ______, *Energy dissipation and self-similar solutions for an unforced inviscid dyadic model*, Trans. Amer. Math. Soc. **363** (2011), 1925–1946.
7. David Barbato, Francesco Morandin, and Marco Romito, *Smooth solutions for the dyadic model*, Nonlinearity **24** (2011), no. 11, 3083–3097.
8. Dirk Blömker, Franco Flandoli, and Marco Romito, *Markovianity and ergodicity for a surface growth PDE*, Ann. Probab. **37** (2009), no. 1, 275–313. `[MR 2489166]`
9. Zdzislaw Brzeźniak, Franco Flandoli, Misha Neklyudov, and Boguslaw Zegarliński, *Conservative interacting particles system with anomalous rate of ergodicity*, 2010, arXiv:1012.0582.
10. Alexey Cheskidov, *Blow-up in finite time for the dyadic model of the Navier-Stokes equations*, Trans. Amer. Math. Soc. **360** (2008), no. 10, 5101–5120.
11. Alexey Cheskidov and Susan Friedlander, *The vanishing viscosity limit for a dyadic model*, Phys. D **238** (2009), no. 8, 783–787.
12. Alexey Cheskidov, Susan Friedlander, and Nataša Pavlović, *Inviscid dyadic model of turbulence: the fixed point and Onsager's conjecture*, J. Math. Phys. **48** (2007), no. 6, 065503, 16.

13. Giuseppe Da Prato and Arnaud Debussche, *Ergodicity for the 3D stochastic Navier-Stokes equations*, J. Math. Pures Appl. (9) **82** (2003), no. 8, 877–947.
14. ______, *On the martingale problem associated to the 2D and 3D stochastic Navier-Stokes equations*, Atti Accad. Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **19** (2008), no. 3, 247–264.
15. Giuseppe Da Prato and Jerzy Zabczyk, *Stochastic equations in infinite dimensions*, Encyclopedia of Mathematics and its Applications, vol. 44, Cambridge University Press, Cambridge, 1992.
16. A. de Bouard and A. Debussche, *On the effect of a noise on the solutions of the focusing supercritical nonlinear Schrödinger equation*, Probab. Theory Related Fields **123** (2002), no. 1, 76–96. `[MR 1906438]`
17. Anne de Bouard and Arnaud Debussche, *Blow-up for the stochastic nonlinear Schrödinger equation with multiplicative noise*, Ann. Probab. **33** (2005), no. 3, 1078–1110. `[MR 2135313]`
18. Arnaud Debussche, *Ergodicity results for the stochastic Navier–Stokes equations: an introduction*, Topics in mathematical fluid–mechanics, Lecture Notes in Math., Springer, Berlin, to appear, Lectures given at the C.I.M.E.–E.M.S. Summer School in applied mathematics held in Cetraro, September 6–11, 2010, Edited by Franco Flandoli and Hugo Beirao da Veiga.
19. Arnaud Debussche and Cyril Odasso, *Markov solutions for the 3D stochastic Navier-Stokes equations with state dependent noise*, J. Evol. Equ. **6** (2006), no. 2, 305–324.
20. John P. Dirkse, *An absorption probability for the Ornstein-Uhlenbeck process*, J. Appl. Probability **12** (1975), no. 3, 595–599.
21. M. Dozzi and J. A. Lopez, *Finite-time blowup and existence of global positive solutions of a semilinear SPDE*, 2009, arXiv:0908.3364.
22. C. L. Fefferman, *Existence and smoothness of the navier-stokes equation*, The millennium prize problems, Clay Math. Inst., Cambridge, MA, 2006, pp. 57–67.
23. Julian Fernández Bonder and Pablo Groisman, *Time-space white noise eliminates global solutions in reaction-diffusion equations*, Phys. D **238** (2009), no. 2, 209–215. `[MR 2516340]`
24. Franco Flandoli, *An introduction to 3D stochastic fluid dynamics*, SPDE in hydrodynamic: recent progress and prospects, Lecture Notes in Math., vol. 1942, Springer, Berlin, 2008, Lectures given at the C.I.M.E. Summer School held in Cetraro, August 29–September 3, 2005, Edited by Giuseppe Da Prato and Michael Röckner, pp. 51–150.
25. ______, *Random perturbation of PDEs and fluid dynamic model*, Lecture Notes in Mathematics, vol. 2015, Springer, Berlin, 2011, École d'Été de Probabilités de Saint-Flour XL.
26. Franco Flandoli and Marco Romito, *Markov selections and their regularity for the three-dimensional stochastic Navier-Stokes equations*, C. R. Math. Acad. Sci. Paris **343** (2006), no. 1, 47–50. `[MR 2241958]`
27. ______, *Regularity of transition semigroups associated to a 3D stochastic Navier-Stokes equation*, Stochastic differential equations: theory and applications (Peter H. Baxendale and Sergey V. Lototski, eds.), Interdiscip. Math. Sci., vol. 2, World Sci. Publ., Hackensack, NJ, 2007, pp. 263–280. `[MR 2393580]`
28. ______, *Markov selections for the 3D stochastic Navier-Stokes equations*, Probab. Theory Related Fields **140** (2008), no. 3-4, 407–458. `[MR 2365480]`
29. Susan Friedlander and Nataša Pavlović, *Blowup in a three-dimensional vector model for the Euler equations*, Comm. Pure Appl. Math. **57** (2004), no. 6, 705–725.
30. ______, *Remarks concerning modified Navier-Stokes equations*, Discrete Contin. Dyn. Syst. **10** (2004), no. 1-2, 269–288, Partial differential equations and applications.
31. Nobuyuki Ikeda and Shinzo Watanabe, *Stochastic differential equations and diffusion processes*, North-Holland Publishing Co., Amsterdam, Oxford, New York, 1981.

32. Ioannis Karatzas and Steven E. Shreve, *Brownian motion and stochastic calculus*, second ed., Graduate Texts in Mathematics, vol. 113, Springer-Verlag, New York, 1991.
33. Nets Hawk Katz and Nataša Pavlović, *Finite time blow-up for a dyadic model of the Euler equations*, Trans. Amer. Math. Soc. **357** (2005), no. 2, 695–708 (electronic).
34. Alexander Kiselev and Andrej Zlatoš, *On discrete models of the Euler equation*, Int. Math. Res. Not. (2005), no. 38, 2315–2339.
35. Nicolai V. Krylov, *The selection of a Markov process from a Markov system of processes, and the construction of quasidiffusion processes*, Izv. Akad. Nauk SSSR Ser. Mat. **37** (1973), 691–708.
36. Carl Mueller, *The critical parameter for the heat equation with a noise term to blow up in finite time*, Ann. Probab. **28** (2000), no. 4, 1735–1746. `[MR 1813841]`
37. Carl Mueller and Richard Sowers, *Blowup for the heat equation with a noise term*, Probab. Theory Related Fields **97** (1993), no. 3, 287–320. `[MR 1245247]`
38. Cyril Odasso, *Exponential mixing for the 3D stochastic Navier-Stokes equations*, Comm. Math. Phys. **270** (2007), no. 1, 109–139.
39. Marco Romito, *Ergodicity of the finite dimensional approximation of the 3D Navier-Stokes equations forced by a degenerate noise*, J. Statist. Phys. **114** (2004), no. 1-2, 155–177. `[MR 2032128]`
40. ______, *Analysis of equilibrium states of Markov solutions to the 3D Navier-Stokes equations driven by additive noise*, J. Stat. Phys. **131** (2008), no. 3, 415–444. `[MR 2386571]`
41. ______, *An almost sure energy inequality for Markov solutions to the 3D Navier-Stokes equations*, 2009, arXiv:0902.1407, to appear on Quaderni di Matematica.
42. ______, *Critical strong Feller regularity for Markov solutions to the Navier–Stokes equations*, J. Math. Anal. Appl. **384** (2010), 115–129, arXiv:1003.4623.
43. ______, *The martingale problem for Markov solutions to the Navier-Stokes equations*, Seminar on Stochastic Analysis, Random Fields and Applications VI, Progr. Probab., vol. 63, Birkhäuser, Basel, 2011, pp. 227–244.
44. Marco Romito and Lihu Xu, *Ergodicity of the 3D stochastic Navier-Stokes equations driven by mildly degenerate noise*, Stoch. Proc. Appl. **121** (2011), no. 4, 673–700.
45. Armen Shirikyan, *Approximate controllability of three-dimensional Navier-Stokes equations*, Comm. Math. Phys. **266** (2006), no. 1, 123–151.
46. Fabian Waleffe, *On some dyadic models of the Euler equations*, Proc. Amer. Math. Soc. **134** (2006), no. 10, 2913–2922 (electronic).

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PISA, LARGO BRUNO PONTECORVO 5, I–56127 PISA, ITALIA
E-mail address: `romito@dm.unipi.it`
URL: `http://www.dm.unipi.it/pages/romito`